



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, C07H 21/04, C12N 15/09, 15/11, 15/85	A1	(11) International Publication Number: WO 99/40789 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/US98/17908 (22) International Filing Date: 28 August 1998 (28.08.98) (30) Priority Data: 60/074,497 12 February 1998 (12.02.98) US 09/108,006 30 June 1998 (30.06.98) US (71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 1100 Washington Avenue South, Minneapolis, MN 55414-1226 (US). ALBERT EINSTEIN COLLEGE OF MEDICINE OF YESHIVA UNIVERSITY [US/US]; 1300 Morris Park Avenue, Bronx, NY 10461 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): STEER, Clifford, J. [US/US]; 1924 Pinchus Avenue, St. Paul, MN 55116-1339 (US). KREN, Betsy, T. [US/US]; 810 29th Avenue S., Minneapolis, MN 55454 (US). BANDYOPADHYAY, Paramita, T. [IN/US]; 23208 Cornerstone Drive, Yardely, PA 19067 (US). ROY-CHOWDHURY, Jayanta [US/US]; 139 Woodhollow Lane, New Rochelle, NY 10804-3435 (US).	(74) Agents: HANSBURG, Daniel; Kimeragen, Inc., 300 Pheasant Run, Newtown, PA 18940 (US) et al. (81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SI, SZ, UG, ZW). Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: METHODS AND COMPOUNDS FOR THE GENETIC TREATMENT OF HYPERLIPIDEMIA		
(57) Abstract <p>The present invention concerns the introduction of specific alterations in the genes that encode three apolipoproteins, Apo A1, Apo B and Apo E. The alterations in Apo A1 introduce a cysteine residue so the disulfide cross-linked Apo A1 homodimers and Apo A1/A2 heterodimers can be formed. The alterations in Apo B introduce stop codons or frame shift mutations that cause the production of a truncated Apo B protein. The alterations in Apo E introduce specific point mutations that have been identified as protective. The production in the liver of a subject of these altered proteins reduces the risk of the subjects developing atherosclerosis. In one embodiment the genetic alterations are introduced by use of chimeric, mixed RNA/DNA, duplex oligonucleotides.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS AND COMPOUNDS FOR THE GENETIC TREATMENT OF HYPERLIPIDEMIA

This application claims benefit of the priority of application Serial No. 60/074,497, filed February 12, 1998 and application Serial No. 09/108,006 filed June 30, 1998.

1. FIELD OF THE INVENTION

The invention concerns methods and compositions for the use of recombinagenic oligonucleobases *in vivo* for the correction of disease causing genetic defects and the prevention of disease by introducing genetic modifications into the genes that encode Apolipoprotein B (Apo B), Apolipoprotein E (Apo E) and Apolipoprotein A1 (Apo A1).

2. BACKGROUND TO THE INVENTION

2.1 THE USE OF CHIMERIC MUTATIONAL VECTORS TO EFFECT GENETIC CHANGES IN CULTURED CELLS

The inclusion of a publication or patent application in this specification is not an admission that the publication or the invention, if any, of the application occurred prior to the present invention or resulted from the conception of a person other than the present inventors.

The published examples of recombinagenic oligonucleobases are termed Chimeric Mutational Vectors (CMV) or chimeraplasts because they contain both 2'-O-modified ribonucleotides and deoxyribonucleotides.

An oligonucleotide having complementary deoxyribonucleotides and ribonucleotides and containing a sequence homologous to a fragment of the bacteriophage M13mp19, was described in Kmiec, E.B., et al., November 1994, Mol. and Cell. Biol. **14**, 7163-7172. The oligonucleotide had a single contiguous segment of ribonucleotides. Kmiec et al. showed that the oligonucleotide was a substrate for the REC2 homologous pairing enzyme from *Ustilago maydis*.

Patent publication WO 95/15972, published June 15, 1995, and counterpart U.S. Patent No. 5,565,350 (the '350 patent) described duplex CMV for the introduction of genetic changes in eukaryotic cells. Examples in a *Ustilago maydis* gene and in the

murine ras gene were reported. The latter example was designed to introduce a transforming mutation into the ras gene so that the successful mutation of the ras gene in NIH 3T3 cells would cause the growth in soft agar of a colony of cells ("transformation"). The '350 patent reported that the maximum rate of transformation of NIH 3T3 was less than 0.1 %, i.e., about 100 transformants per 10^6 cells exposed to the ras duplex CMV. In the *Ustilago maydis* system, the rate of transformants was about 600 per 10^6 . A chimeric vector designed to introduce a mutation into a human bcl-2 gene was described in Kmiec, E.B., February 1996, Seminars in Oncology 23, 188.

A duplex CMV designed to repair the mutation in codon 12 of K-ras was described in Kmiec, E.B., December 1995, Advanced Drug Delivery Reviews 17, 333-40. The duplex CMV was tested in Capan 2, a cell line derived from a human pancreatic adenocarcinoma, using LIPOFECTIN™ to introduce the duplex CMV into the Capan 2 cells. Twenty four hours after the duplex CMV was introduced, the cells were harvested and genomic DNA was extracted; a fragment containing codon 12 of K-ras was amplified by PCR and the rate of conversion estimated by hybridization with allele specific probes. The rate of repair was reported to be approximately 18%.

A duplex CMV designed to repair a mutation in the gene encoding liver/bone/kidney type alkaline phosphatase was reported in Yoon, K., et al., March 1996, Proc. Natl. Acad. Sci. 93, 2071. The alkaline phosphatase gene was transiently introduced into CHO cells by a plasmid. Six hours later the duplex CMV was introduced. The plasmid was recovered at 24 hours after introduction of the duplex CMV and analyzed. The results showed that approximately 30 to 38% of the alkaline phosphatase genes were repaired by the duplex CMV.

WO 97/41411 and counterpart United States Patent No. 5,760,012 to E.B. Kmiec, A. Cole-Strauss and K. Yoon, and the publication Cole-Strauss, A., et al., September 1996, SCIENCE 273, 1386 disclose duplex CMV that are used in the treatment of genetic diseases of hematopoietic cells, e.g., Sickle Cell Disease, Thalassemia and Gaucher Disease. United States Patent Application Serial No. 08/664,487, filed June 17, 1996, by E.B. Kmiec describes duplex CMV having non-natural nucleotides for use in specific, site-directed mutagenesis. The duplex CMV described in the applications and certain of the publications of Kmiec and his colleagues contain a central segment of

DNA:DNA homoduplex and flanking segments of RNA:DNA hybrid-duplex or 2'-OMe-RNA:DNA hybrid-duplex.

The work of Kmiec and his colleagues concerned cells that are mitotically active, i.e., proliferating cells, at the time they are exposed to CMV. Kmiec and colleagues used a CMV/liposomal macromolecular carrier complex in which the CMV were mixed with a pre-formed liposome or lipid vesicle. In such a complex the CMV are believed to adhere to the surface of the liposome.

Kren et al., June 1997, *Hepatology* **25**, 1462-1468, reported the successful use of a CMV in non-replicating, primary tissue-cultured rat hepatocytes to mutate the coagulation factor IX gene. Kren et al., March 1998, *Nature Medicine* **4**, 285 reported the use of a CMV *in vivo* to introduce a genetic defect in the same gene.

2.2 THE USE OF A POLYETHYLENIMINE MACROMOLECULAR CARRIER FOR *IN VIVO* AND *IN VITRO* TRANSFECTION

Branched chain polyethylenimine has been used as a carrier to introduce nucleic acids into eukaryotic cells both *in vivo* and *in vitro*. Boussif, O., et al., 1995, *Proc. Natl. Acad. Sci.* **92**, 7297; Abdallah, B. et al., 1996, *Human Gene Therapy* **7**, 1947. Boletta, A., et al., 1997, **8**, 1243-1251. The *in vitro* use of galactosylated polyethylenimine to introduce DNA into cultured HepG2 hepatocarcinoma cell lines is reported by Zanta, et al., October 1, 1997, *Bioconjugate Chemistry* **8**, 839-844. The coupling of a protein ligand, transferrin, to polyethylenimine and its use to introduce a test gene into cultured cells by use of the transferrin receptor is described in Kircheis, R., et al., 1997, *Gene Therapy* **4**, 409-4-18. Branched chain polyethylenimines contain secondary and tertiary amino groups having a broad range of pK's and, consequently these polyethylenimines have a substantial buffering capacity at a pH where polylysine has little or no capacity, i.e., less than about 8. Tang, M.K., & Szoka, F.C., 1997, *Gene Therapy* **4**, 823-832. The use of branched chain polyalkanylimines, including polyethylenimine as carriers for the introduction of nucleic acids into cells is described in WO 96/02655 to J-P. Behr et al.

The successful *in vivo* and *in vitro* use of linear polyethylenimine to transfect a gene is reported by Ferrari, S., et al., 1997, *Gene Therapy* **4**, 1100-1106. Compositions

comprising a linear polyalkanylimine and a nucleic acid as disclosed in patent publication WO 93/20090 to S. Stein et al.

2.3 THE USE OF A LIPOSOMAL CARRIER FOR *IN VIVO* TRANSFECTION

The use of liposomes or lipid vesicles to introduce DNA encoding a foreign protein into cells has been described. The most frequently used techniques adhere the DNA to the surface of a positively charged liposome, rather than encapsulating the DNA, although encapsulated DNA techniques were known. United States Patent Nos. 4,235,871 and 4,394,448 are relevant. The field is reviewed by Smith, J.G., et al., 1993, *Biochim. Biophys. Acta* **1154**, 327-340 and Staubinger, R.M., et al., 1987, *Methods in Enzymology* **185**, 512. The use of DOTAP, a cationic lipid in a liposome to transfect hepatic cells *in vivo* is described in Fabrega, A.J., et al., 1996, *Transplantation* **62**, 1866-1871. The use of cationic lipid-containing liposomes to transfect a variety of cells of adult mice is described in Zhu, N., et al., 1993, *Science* **261**, 209. The use of phosphatidylserine containing lipids to form DNA encapsulating liposomes for transfection is described in Fraley, R., et al., 1981, *Biochemistry* **20**, 6978-87.

2.4 THE USE OF THE ASIALOGLYCOPROTEIN RECEPTOR FOR HEPATOCELLULAR SPECIFIC TRANSFECTION

United States Patent Nos. 5,166,320 and 5,635,383 disclose the transfection of hepatocytes by forming a complex of a DNA, a polycationic macromolecular carrier and a ligand for the asialoglycoprotein receptor. In one embodiment, the macromolecular carrier was polylysine. The use of a lactosylcerebroside containing liposome to transfect a hepatocyte *in vivo* is described by Nandi, P.K., et al., 1986, *J. Biol. Chem.* **261**, 16722-16722. The use of asialofetuin-labeled liposomes to transfect liver cells with a reporter plasmid is described in Hara et al., 1995, *Gene Therapy* **2**, 764-788. The use of galactosylated polyethyleneimine to transfect cultured hepatocytes is described in Zanta M-A., et al. abst. pub. Oct. 1, 1997, *Bioconjugate Chem.*, **8**, 839-844.

2.5 APO B100, APO B48 AND THE REDUCTION OF SERUM LDL

Hepatic and Intestinal Lipoprotein Secretion: Both the liver and the intestines make and export lipoproteins for the transport of lipids. The lipoproteins are termed very low density lipoproteins (VLDL) and chylomicrons, respectively. VLDL and chylomicrons differ in size and in their major protein components. The major protein of VLDL is Apo B100, consisting of 4536 amino acids; the major protein of chylomicrons is Apo B48, which consists of the N-terminal 2152 amino acids of Apo B100. Apo B48 and Apo B100 are encoded by a single gene, the transcript of which is modified at nucleotide 6666 (codon 2179) by a sequence specific cytidine deaminase, termed apolipoprotein B mRNA editing enzyme (APOBE). The action of this enzyme converts a C to U and results in a stop codon.

Both VLDL, which contain Apo B100, and chylomicrons, which contain Apo B48 transport triglycerides in the vascular system to a delivery site. However, after triglyceride hydrolysis and delivery VLDL are transformed into LDL, while chylomicrons are not. High levels of circulating LDL *per se* and a high LDL:HDL ratio increase the risk of arterial atherosclerosis. Hence, it has been suggested that increasing the ratio of Apo B48 to Apo B100 would have a beneficial effect.

In many species of mammals, e.g., rats and mice, a high percentage of the lipid secretions of both liver and intestine contain Apo B48. Such species have markedly lower ratios of LDL:HDL. Greve J., et al., 1995, Proc. Zool. Soc., Calcutta, **47**, 93-100. In others, such as humans and rabbits, hepatocytes lack APOBE and the hepatocytes consequently produce only VLDL.

One strategy to reduce the atherosclerosis in humans has been to introduce the gene for the catalytic component of the apolipoprotein B editing enzyme (APOBEC-1) under the control of a constitutive promoter to convert Apo B100 transcripts into Apo B48 transcripts. The transient expression of APOBEC-1 in the hepatocytes of normal and genetically hyperlipidemic Watanabe rabbit does cause a transient reduction in the levels of LDL. Greeve, J., et al., 1996, J. Lipid Res. **37**, 2001-17. However, the uncontrolled production of APOBEC-1 is mutagenic and may cause hepatocellular hyperplasia and hepatocellular carcinoma. Yamanaka, S., et al., 1995, Proc. Natl. Acad. Sci. **92**, 8483-8487.

Individuals who are homozygous or mixed heterozygotes for genes encoding truncated Apo B100 have been observed. Malloy et al., 1981, J. Clin. Invest. **67**, 1441; Hardman, D.A., et al., 1991, J. Clin. Invest. **88**, 1722. These individuals have low or absent LDL. For example, deletion of nucleotides 5391-5394 results in a frame shift mutation and a shortened Apo B (B37). These patients are most often asymptomatic. Steinberg, D., et al., 1979, J. Clin. Invest. **64**, 292; Young, S.G., et al., 1988, Science **241**, 591; Young, S.G., 1987, J. Clin. Invest. **79**, 1831. Reviewed Linton, M.F., 1993, J. Lipid. Res. **34**, 521; Kane, J.P. & Havel, R.J., 1995, Chapt. 57, THE METABOLIC BASIS OF INHERITED DISEASE, ed. Scriver et al. (McGraw Hill, New York). Similarly, as many as 1 in every 3,000 persons has a serum cholesterol level of 100 mg/dl or less because the individual is heterozygous for a truncated Apo B gene. *Ibid.*, p. 1866.

Truncations that result in an Apo B that are shorter than Apo B 31 do not circulate. Truncated Apo B 86, 87 and 90 have been observed. Apo B 86 and Apo B 87, are not associated with LDL while Apo B 90 is. Each mutation is associated with hypobetalipoproteinemia. Linton, M.L., et al., 1990, Clin. Res. **38**, 286A (abstr.); Tennyson, G.E., et al., 1990, Clin. Res. **38**, 482A (abstr.); Kruhl, E.S., et al., 1989, Arteriosclerosis **9**, 856.

2.6 APO E POLYMORPHISM AND TYPE III HYPERLIPIDEMIA

Apolipoprotein E is the major ligand for the LDL receptor for lipoproteins that contain Apo B48. There are three allelic forms of human Apo E that differ from each other by one or two amino acids: Apo E2 (Cys¹¹² Cys¹⁵⁸); Apo E3 (Cys¹¹² Arg¹⁵⁸); and Apo E4 (Arg¹¹² Arg¹⁵⁸). There is considerable geographical variation in the prevalences of the alleles. Excluding Africa, E2 ranges between 4% and 12 %, E3 between 70% and 85% and E4 between 7.5 and 25%. In the Sudan, the prevalences are 8.1%, 61.9% and 29.1%, respectively. Mahley, R.W. & Rall, S.C., Jr., 1995, Chapt. 61, THE METABOLIC BASIS OF INHERITED DISEASE, ed. Scriver et al. (McGraw Hill, New York). Thus approximately 1% of the North American and European population are Apo E 2/2 homozygotes. Of these homozygotes approximately between 2% and 10% display type III hyperlipidemia. Paradoxically, however, Apo E 2/2 homozygotes that have not

developed overt Type III hyperlipidemia display lower than average LDL associated cholesterol. Davignon, J., 1988, *Arteriosclerosis* **8**, 1.

The E4 allele is also associated with increased incidence of a major disease, Alzheimer's Disease, and with increased risk of coronary artery disease. Roses, A.D., 1996, *Ann. NY Acad. Sci.* **802**, 50-57; Okumoto, K., & Fujiki, Y., 1997, *Nature Genetics* **17**, 263; Kuusi, T., et al., 1989, *Arteriosclerosis* **9**, 237. A polymorphism in the region 491 nt 5' to the transcription start site of the Apo E gene is also an independently associated with increased risk of Alzheimer's disease. Individuals homozygous for the -491-A genotype have an increased risk of Alzheimer's, while individuals homozygous or heterozygous for the -491 T genotype have no increased risk. Bullido, M.J., 1998, et al., *Nature Genetics* **18**, 69-71.

The E2 allele in most individuals is associated with the lowest levels of serum cholesterol and LDL. However, about 5% of E2/E2 homozygous persons who are subject to environmental or genetic stress develop type III hyperlipidemia. The most common stressors are hypothyroidism, untreated diabetes mellitus, alcoholism and marked weight gain. Removal of the stressor usually results in control of the hyperlipidemia. Rare patients with type III hyperlipidemia have mutant Apo E genes. Mahley & Rall, *ibid.* Table 61-5.

2.7

APO A1 AND HDL

High density lipoproteins (HDL) transport cholesterol and phospholipids from peripheral extrahepatic locations to the liver. In particular, HDL are believed to remove lipid deposits from vascular endothelial cells and that the observed negative correlation between the levels of HDL-cholesterol (HDL-C) and coronary artery disease is due to this function. Eisenberg, S., 1984, *J. Lipid Research* **25**, 1017; Gordon, D.J., et al., 1986, *Circulation* **74**, 1217. HDL are secreted by the liver and intestines as nascent HDL particles containing four molecules of apo A1, which is a 243 amino acid protein. The nascent HDL physically attract free cholesterol from cell membranes and/or other lipoproteins. The resulting particle contains apo A1, phospholipid, and cholesterol. Such particles are substrates for lecithin:cholesterol acyltransferase (LCAT), which esterifies the free cholesterol to cholesterol esters. The presence of the more hydrophobic cholesterol

esters transforms the nascent HDL initially to a more stable mature HDL3 and subsequently HDL3 particle.

Cholesterol ester transfer protein removes the esterified cholesterol from the HDL3 and HDL2 particles and transfers them into LDL and thence into hepatocytes through the LDL receptors.

Two mutations of apo A1 have been discovered wherein the levels of HDL are depressed. The mutations are missense mutations that replace an arginine with a cysteine amino acid. Weisgraber, K.H., et al., 1983, J. Biol. Chem. **258**, 2508 (apo A1 milano (Arg-Cys)¹⁷³); Bruckert, E., et al., 1997, Atherosclerosis **128**, 121 (apo A1 R151C (Arg-Cys)¹⁵¹). The mutation is dominant, i.e., HDL levels are depressed in affected individuals who are heterozygous for the mutation. An explanation for this is that the mutant apo A1 proteins form cystine linked heterodimers with wild apo A2 molecules, which are a minor apolipoprotein found in HDL, as well as homodimers.

Surprising even though the lipid profile, i.e., low HDL and elevated triglycerides, is one that ordinarily is associated with accelerated atherosclerosis, the mutations are not associated with atherosclerosis, but, rather, are believed to be protective of atherosclerosis.

The mechanism of protection is not established. A large (N = 33) retrospective epidemiologic study of apo A1 milano carriers shows that there is a low incidence of coronary artery disease in these individuals, however, the significance of this fact must be viewed in the context of similar low incidence among non-affected residents of the village Limone sul Garda. Gualandri, V., et al., 1985, Am. J. Hum. Gen. **37**, 1083. *In vitro* studies indicate that the mutant apo A1 is not more effective in recruiting plasma membrane cholesterol. Bielicki, J.K., et al., 1997, Arterioscler Thromb Vasc Biol **17**, 1637. There is direct evidence that the administration of exogenous apo A1 milano, but not wild type apo A1, in the form of a complex with phospholipids, is protective in an accepted model system of atherosclerosis. Ameli, S. et al., 1994, Circulation **90**, 1935; Shah, P.K., et al., 1998, Circulation **97**, 780.

3. SUMMARY OF THE INVENTION

The present invention concerns methods of treatment and/or prophylaxis which consists of the introduction of specific genetic alterations in genes of a subject individual. In one embodiment, the specific genetic alteration blocks the synthesis of Apo B100 and thereby reduces the level of LDL cholesterol. In an alternative embodiment, the specific alteration converts an Apo E4 allele to an Apo E3 or Apo E2 allele, which is associated with decreased risk of atherosclerosis and Alzheimer's Disease. In further alternative embodiments, the invention concerns the correction of inherited genetic defects in the genes of hepatocytes of individuals having a disease caused by such defects.

The present invention further comprises a method of treating and/or preventing atherosclerosis by causing mutations in the genes encoding apo A1 and compounds that are useful for the introduction of such mutations. The mutations useful for the practice of the invention are mutations that insert a cysteine residue that forms a cystine and results in the formation of homodimers and apo A2-containing heterodimers.

The embodiments of the invention can be practiced using any oligonucleotide or analog or derivative thereof, now known or hereafter developed, that can cause specific genetic alterations in the genome of the hepatocytes of the subject individual (hereafter a "recombinagenic oligonucleobase"), for example a chimeric mutational vector (CMV) as, for example, described in United States patent No. 5,565,350, No. 5,731,181, and No. 5,760,012. Alternatively, the recombinagenic oligonucleobase can be a heteroduplex mutational vector or a non-chimeric mutational vector as described in U.S. patent application No. 09/078,063 and No. 09/078,064, filed May 12, 1998, each of which are hereby incorporated by reference.

In a preferred embodiment the recombinagenic oligonucleobase is complexed with a macromolecular carrier to which is attached a specific ligand. The ligand is selected to bind to a cell-surface receptor that is internalized into hepatocytes through clathrin-coated pits into endosomes. The cell surface receptors that bind such ligands are termed herein "clathrin-coated pit receptors". Examples of hepatic clathrin-coated pit receptors include the low density lipoprotein (LDL) receptor and the asialoglycoprotein receptor.

In specific embodiments the macromolecular carrier can be 1) an aqueous-cored lipid vesicle of between 25 nm and 400 nm diameter, wherein the aqueous core contains the CMV; 2) a lipid nanosphere of between 25 nm and 400 nm diameter, having a lipid core, wherein the lipid core contains a lipophilic salt of the CMV; or 3) a polycationic salt of the CMV. Examples of polycations for such salts include polyethylenimine, polylysine and histone H1. In one embodiment the polycation is a linear polyethylenimine (PEI) salt having a mass average molecular weight greater than 500 daltons and less than 1.3 Md. Alternatively the polycation can be a branched-chain polyethylenimine.

4. BRIEF DESCRIPTION OF THE FIGURE

Figure 1 is a schematic of one embodiment of CMV useful in the invention.

Figures 2A-2C show the genomic sequence of human APO E gene with translation of exons. Introns are in lower case and exons are in upper case.

Figure 3A-3G shows the sequence of the human APO A1 gene (SEQ ID No. 59)

5. DEFINITIONS

The invention is to be understood in accordance with the following definitions.

An oligonucleobase is a polymer of nucleobases, which polymer can hybridize by Watson-Crick base pairing to a DNA having the complementary sequence.

Nucleobases comprise a base, which is a purine, pyrimidine, or a derivative or analog thereof. Nucleobases include peptide nucleobases, the subunits of peptide nucleic acids, and morpholine nucleobases as well as nucleosides and nucleotides.

Nucleosides are nucleobases that contain a pentosefuranosyl moiety, e.g., an optionally substituted riboside or 2'-deoxyriboside. Nucleosides can be linked by one of several linkage moieties, which may or may not contain a phosphorus. Nucleosides that are linked by unsubstituted phosphodiester linkages are termed nucleotides.

An oligonucleobase chain has a single 5' and 3' terminus, which are the ultimate nucleobases of the polymer. A particular oligonucleobase chain can contain nucleobases of all types. An oligonucleobase compound is a compound comprising one or more oligonucleobase chains that are complementary and hybridized by Watson-Crick base pairing. Nucleobases are either deoxyribo-type or ribo-type. Ribo-type nucleobases are

pentosefuranosyl containing nucleobases wherein the 2' carbon is a methylene substituted with a hydroxyl, alkyloxy or halogen. Deoxyribo-type nucleobases are nucleobases other than ribo-type nucleobases and include all nucleobases that do not contain a pentosefuranosyl moiety.

An oligonucleobase strand generically includes both oligonucleobase chains and segments or regions of oligonucleobase chains. An oligonucleobase strand has a 3' end and a 5' end. When a oligonucleobase strand is coextensive with a chain, the 3' and 5' ends of the strand are also 3' and 5' termini of the chain.

A region is a portion of an oligonucleobase, the sequence of which is derived from some particular source, e.g., a CMV having a region of at least 15 nucleotides having the sequence of a fragment of the human β -globin gene. A segment is a portion of a CMV having some characteristic structural feature. A given segment or a given region can contain both 2'-deoxynucleotides and ribonucleotides. However, a ribo-type segment or a 2'-deoxyribo-type segment contain only ribo-type and 2'-deoxyribo-type nucleobases, respectively.

6. DETAILED DESCRIPTION OF THE INVENTION

6.1 THE STRUCTURE OF THE CHIMERIC MUTATIONAL VECTOR

The Chimeric Mutational Vectors (CMV) are comprised of oligonucleobases, i.e., polymers of nucleobases, which polymers form Watson-Crick base pairs of purines and pyrimidines (hybridize), to DNA having the appropriate sequence. Each CMV is divided into a first and a second strand of at least 15 nucleobases each that are complementary to each other. The strands can be, but need not be, covalently linked. Nucleobases contain a base, which is either a purine or a pyrimidine or analog or derivative thereof. There are two types of nucleobases. Ribo-type nucleobases are ribonucleosides having a 2'-hydroxyl, substituted 2'-hydroxyl or 2'-halo-substituted ribose. All nucleobases other than ribo-type nucleobases are deoxyribo-type nucleobases. Thus, deoxy-type nucleobases include peptide nucleobases. As used herein, only a recombinagenic oligonucleobase that contains at least three contiguous ribo-type nucleobases that are hybridized to deoxyribo-type nucleobases are considered CMV.

The sequence of the first and second strands consists of at least two regions that are homologous to the target gene, i.e., have the same sequence as fragments of the target gene, and one or more regions (the "mutator regions") that differ from the target gene and introduce the genetic change into the target gene. The mutator region is located between homologous regions. In certain embodiments of the invention, each of the flanking homologous regions contains a ribo-type segment of at least three ribo-type nucleobases, that form a hybrid duplex, preferably at least six ribo-type nucleobases and more preferably at least ten ribo-type nucleobases in length, but not more than 25 and preferably not more than 20, more preferably not more than 15 ribo-type nucleobases. The hybrid-duplex-forming ribo-type oligonucleobase segments need not be adjacent to the mutator region. In certain embodiments of the invention the ribo-type oligonucleobase segments are separated from the mutator region by a portion of the homologous region comprising deoxyribo-type nucleobases. In these embodiments the mutator region is also composed of deoxyribo-type nucleobases. Accordingly, the mutator region and a portion of one or both homologous regions form an intervening segment of homo-duplex, which separates the two segments of hybrid-duplex.

The total length of all homologous regions is preferably at least 16 nucleobases and is more preferably from about 20 nucleobases to about 60 nucleobases in length.

Preferably, the mutator region consists of 20 or fewer bases, more preferably 6 or fewer bases and most preferably 3 or fewer bases. The mutator region can be of a length different than the length of the sequence that separates the regions of the target gene homology with the homologous regions of the CMV so that an insertion or deletion of the target gene results. When the CMV is used to introduce a deletion in the target gene there is no base identifiable as within the mutator region. Rather, the mutation is effected by the juxtaposition of the two homologous regions that are separated in the target gene. For the purposes of the invention, the length of the mutator region of a CMV that introduces a deletion in the target gene is deemed to be the length of the deletion. In one embodiment the mutator region is a deletion of from 6 to 1 bases or more preferably from 3 to 1 bases. Multiple separated mutations can be introduced by a single CMV, in which case there are multiple mutator regions in the same CMV. Alternatively multiple CMV can be used simultaneously to introduce multiple genetic changes in a single gene

or, alternatively to introduce genetic changes in multiple genes of the same cell. Herein the mutator region is also termed the heterologous region.

In one embodiment the CMV is a single oligonucleobase chain of between 40 and 100 nucleobases. In an alternative embodiment, the CMV comprises a first and a second oligonucleobase chain, each of between 20 and 100 bases; wherein the first chain comprises the first strand and the second chain comprises the second strand. The first and second chains can be linked covalently by other than nucleobases or, alternatively, can be associated only by Watson-Crick base pairings. In an alternative embodiment the CMV is a first strand which is a single oligonucleobase chain and a second strand, complementary to the first which consists of two oligonucleobase chains, which are linked to the first strand chain by linkers. The combined length of the two chains of the second strand is the length of the first strand.

Linkers: Covalent linkage of the first and second strands can be made by oligo-alkanediols such as polyethyleneglycol, poly-1,3-propanediol or poly-1,4-butanediol. The length of various linkers suitable for connecting two hybridized nucleic acid strands is understood by those skilled in the art. A polyethylene glycol linker having from six to three ethylene units and terminal phosphoryl moieties is suitable. Durand, M. et al., 1990, *Nucleic Acid Research* **18**, 6353; Ma, M. Y-X., et al., 1993, *Nucleic Acids Res.* **21**, 2585-2589. A preferred alternative linker is bis-phosphorylpropyl-trans-4,4'-stilbenedicarboxamide. Letsinger, R.L., et alia, 1994, *J. Am. Chem. Soc.* **116**, 811-812; Letsinger, R.L. et alia, 1995, *J. Am. Chem. Soc.* **117**, 7323-7328, which are hereby incorporated by reference. Such linkers can be inserted into the CMV using conventional solid phase synthesis. Alternatively, the strands of the CMV can be separately synthesized and then hybridized and the interstrand linkage formed using a thiophoryl-containing stilbenedicarboxamide as described in patent publication WO 97/05284, February 13, 1997, to Letsinger R.L. et alia.

In a further alternative embodiment the linker can be a single strand oligonucleobase comprised of nuclease resistant nucleobases, e.g., a 2'-O-methyl, 2'-O-allyl or 2'-F-ribonucleotides. The tetranucleotide sequences TTTT, UUUU and UUCC and the trinucleotide sequences TTT, UUU, or UCG are particularly preferred nucleotide linkers.

Nucleotides: In an alternative embodiment the invention can be practiced using CMV comprising deoxynucleotides or deoxynucleosides and 2'-O substituted ribonucleotides or ribonucleosides. Suitable substituents include the substituents taught by the Kmiec Application, C₁₋₆ alkane. Alternative substituents include the substituents taught by U.S. Patent No. 5,334,711 (Sproat) and the substituents taught by patent publications EP 629 387 and EP 679657 (collectively, the Martin Applications), which are hereby incorporated by reference. As used herein a 2' fluoro, chloro or bromo derivative of a ribonucleotide or a ribonucleotide having a substituted 2'-O as described in the Martin Applications or Sproat is termed a "2'-Substituted Ribonucleotide." Particular preferred embodiments of 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-propyloxy, 2'-allyloxy, 2'-hydroxylethyloxy, 2'-methoxyethyloxy, 2'-fluoropropyloxy and 2'-trifluoropropyloxy substituted ribonucleotides. In more preferred embodiments the 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-methoxyethyloxy, and 2'-allyloxy substituted nucleotides.

2'-Substituted Ribonucleosides are defined analogously. Particular preferred embodiments of 2'-Substituted Ribonucleosides are 2'-fluoro, 2'-methoxy, 2'-propyloxy, 2'-allyloxy, 2'-hydroxylethyloxy, 2'-methoxyethyloxy, 2'-fluoropropyloxy and 2'-trifluoropropyloxy substituted ribonucleotides. In more preferred embodiment on the 2'-Substituted Ribonucleosides are 2'-fluoro, 2'-methoxy, 2'-methoxyethyloxy, and 2'-allyloxy substituted nucleotides.

The term "nuclease resistant ribonucleoside" encompasses 2'-Substituted Ribonucleosides, including 2'-Substituted Ribonucleotides and also all 2'-hydroxyl ribonucleosides other than ribonucleotides. In a preferred embodiment, the CMV preferably includes at least three and more preferably six nuclease resistant ribonucleosides. In one preferred embodiment the CMV contains no nuclease sensitive ribonucleosides. In an alternative preferred embodiment, every other ribonucleoside is nuclease resistant. Certain 2'-blocking groups can be more readily synthesized for purines or pyrimidines. In one embodiment of the CMV only the ribonucleoside purines or only the ribonucleoside pyrimidines are nuclease resistant.

Recombinagenic oligonucleobases, including non-chimeric mutational oligonucleobases and improved CMV and their use in eukaryotic cells and cell-free

systems are described in U.S. patent applications Serial No. 09/078,063, filed May 12, 1998, and Serial No. 09/078,064, filed May 12, 1998, which are each hereby incorporated in their entirety. These mutational oligonucleobases can be used in the same manner as the CMV described in this application.

6.2 THE GENE-SPECIFIC STRUCTURE OF THE CHIMERIC MUTATIONAL VECTOR

Figure 1 shows a diagram of a CMV according to one embodiment of the invention. In the Figure segments "a" and "c-e" are target gene specific segments of the CMV. The sequence of segment "a" and "c-e" are complements of each other. The sequence of segments "f" and "h" are also complements of each other but are unrelated to the specific target gene and are selected merely to ensure the stability of hybridization in order to protect the 3' and 5' ends. Additional protection of the 3' and 5' ends can be accomplished by making the 5' and 3' most internucleotide bonds a phosphorothioate, phosphonate or any other nuclease resistant bond. The sequence of segments "f" and "h" can be 5'-GCCGC-3' or permutations thereof. Segments "g" and "b" can be any linker that covalently connects the two strands, e.g., four unpaired nucleotides or an alkoxy oligomer such as polyethylene glycol. When segments "g" and "b" are composed of other than nucleobases, then segments "a", "c-f" and "h" are each an oligonucleobase chain.

The ribo-type nucleobase segments are segments "c" and "e," which form hybrid-duplexes by Watson-Crick base pairing to the complementary portions of segment "a." The segment "a" can have the sequence of either the coding or non-coding strand of the gene.

Table I contains SEQ ID No. 4 - No. 21 and Table III contains SEQ ID No. 22-25 and 54-58, which are examples of the sequences that can be used to practice the invention. The mutator region in each case is underlined and in bold. CMV having a segment "a" with a sequence selected from the sequences of Table I can be used to practice the invention. Alternatively, segment "a" may have the sequence of the complement of a sequence of Table I. As used herein, a CMV or other type of recombinagenic oligonucleobase comprises a sequence if either strand of the CMV or recombinagenic oligonucleobase comprises the sequence or comprises a sequence containing ribo-type nucleobases with uracil bases replacing thymine bases. Thus, for

example, a CMV having the sequence 5'-agucuggaugGCTAAgccgcccuca-3' (SEQ ID No. 26) is considered to have the sequence of SEQ ID No: 4, wherein the lower case letters denote ribo-type nucleobases and the UPPER CASE letters denote deoxyribo-type nucleobases.

Subjects can be treated with a recombinagenic oligonucleobase specific for Apo B or Apo E according to the guidance of the Factor IX example below. More particularly the recombinagenic oligonucleobase can be given in divided doses at intervals that permit determining of the phenotypic effect of the dose, i.e., evaluation of the extent of the decline in LDL cholesterol and observation for adverse reactions. A reduction of the subject's fasting LDL serum cholesterol to below the level of the 5th percentile of the age-matched population (80-90 mg/dl) can be used as a therapeutic end point; alternatively reduction of fasting LDL serum cholesterol to below the average age-matched normal value (100-140) can be used. The number and size of the dose(s) can be modified to control the extent of the phenotypic effects. In the event that reversal of the specific genetic changes appear desirable, a recombinagenic oligonucleobase having a sequence appropriate to reverse the specific changes can be administered so that the fraction of unmodified Apo B or Apo E genes can be increased. Modification of the dose size and number and the administration of a reversing recombinagenic oligonucleobase permits the adjustment of the number of altered genes in the subject so that a predetermined amount of the phenotypic change can be effected.

6.2.1 Specific Alterations of the Apo B Gene

SEQ ID No. 1 contains the Apo B amino acid sequence and SEQ ID No. 2 contains the Apo B cDNA sequence.

The level of serum cholesterol and particularly of LDL-associated cholesterol can be reduced in a subject by introducing mutations into the subject's hepatic Apo B genes. The mutation can be any mutation that causes termination of the Apo B translation product between amino acid 1433 (Apo B 31) and amino acid 3974 (Apo B 87). (The amino acid numbering for Apo B in this specification refers to the 4553 amino acid primary translation product, i.e., mature Apo B100 plus the 27 amino acid leader sequence. Mature Apo B 100 consists of 4536 amino acids and mature Apo B 48 consists

of 2152 amino acids.) Preferably the translation product is terminated between amino acids 1841 (Apo B 40) and 2975 (Apo 65). The translation product can be terminated by introducing a frameshift mutation, i.e., by adding or deleting one or two nucleotides from the gene, or by introducing a stop codon (a TAA, TAG or TGA). The preferred stop codon is TAA. To monitor the introduction of the mutation it is preferred to have the mutation introduce or remove a palindromic sequence, which is the substrate of a restriction enzyme.

The sequence of the CMV is selected to have two homologous regions of at least 10 nucleobases and preferably at least 12 nucleobases each with a fragment of the Apo B gene located between nucleotides encoding amino acid 1433 (nt 4425) and 3974 (nt 12,048) and preferably located between the nucleotides encoding amino acids 1841 (nt 5649) and 2975 (nt 9051). In this specification, nt 6666 is the first nucleotide of codon 2180, the nucleotide that is converted by APOBE. In a preferred embodiment, the two homologous regions are separated by a single nucleobase in the sequence of the Apo B gene, where the CMV introduces a base substitution in the Apo B gene. Alternatively, the two homology regions can be adjacent in the Apo B gene and separated by a single or double nucleobase in the CMV, such that a one or two base insertion results from the action of the CMV on the Apo B gene. Alternatively, the homologous regions can be separated in the Apo B gene by one or two nucleotides that are deleted from the sequence of the CMV, such that the action of the CMV results in a one or two base deletion in the gene.

Nucleotides 4425-12,048 of the Apo B cDNA are encoded by exon 26 (nt 4342-11913), exon 27 (nt 11914 - 12028) and exon 28 (nt 12029-12212); see Table I, and GENBANK Accession No. 19828, which is hereby incorporated by reference. When an alteration is to be made at a position 3' of nt 11913, attention must be paid to the exon/intron boundary. Mutations that are located within 10-15 nucleotides of the exon/intron boundary must be identified so that the homology region of the CMV continues with the sequence of the intron and not the exon.

The homologous regions can be each from 10 to about 15 nucleobases in length; the two regions need not be of the same length. The fraction of nucleobases that contain a guanine or cytosine base is a design consideration (the GC fraction). It is preferred that

when the homologous region contains 12 or fewer nucleobases, the GC fraction be at least 33% and preferably at least 50%. When the GC fraction is less than 33% the length of the homologous regions is preferably 13, 14 or 15 nucleobases.

Table I contains 18 exemplary embodiments, SEQ ID No. 4-21 and Table III contains 9 exemplary embodiments, SEQ ID No. 22-25 and 54-58, of CMV sufficient for the practice of the embodiments of the invention described in this section. Suitable CMV can be made using nt 3-23 of SEQ ID No. 4-10, 12, and 16-20. SEQ ID NO. 11 and 13-15 have a lower GC fraction; CMV sufficient for the practice of the invention can be made containing residues 3-25 of SEQ ID NO. 11 and 13-15.

6.2.2 Specific Alterations of the Apo E Gene

In a further embodiment, the invention consists of introducing specific alterations to the Apo E gene. E4 homozygous individuals are at increased risk for atherosclerosis, particularly coronary artery disease, and Alzheimer's disease. Therefore, one embodiment of the present invention is the introduction of the substitution Arg→Cys at residues 112, to convert an E4 allele to an E3 allele, and optionally at residue 158 to convert an E3 or E4 allele into an E2 allele of an Apo E gene of an hepatocyte of a subject. The substitutions can be introduced using an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 22 and No. 23 or complement thereof and more preferably of an oligonucleobase containing SEQ ID No. 22 and No. 23 or complement thereof. In addition, in individuals lacking genetic or environment stressors, the E2 allele results in a lowered LDL level and a decreased risk of atherosclerosis and coronary artery disease. Thus, these risks in an E3/E3 individual can be reduced by introduction of the (Arg→Cys)¹⁵⁸ substitution to convert the individual Apo E genes to E2 alleles.

Apo E2/E2 homozygous individuals who are suffering from Type III hyperlipidemia can be treated by converting E2 alleles to E3 alleles by making a Cys→Arg¹⁵⁸ substitution. Such a substitution can be made using an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 24 or complement thereof and more preferably of an oligonucleobase containing SEQ ID No. 24 or complement thereof.

Independent of the Apo E allele, individuals who are homozygous for -491-A are at increased risk to develop Alzheimer's Disease. Bullido, M.J., 1998, et al., Nature

Genetics **18**, 69-71. These individuals can be advantageously treated with an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 25.

6.2.3 Repair of Mutations of the Apo B and Apo E Gene

SEQ ID No. 3 contains the Apo E genomic DNA sequence.

A further embodiment of the invention concerns the use of CMV to repair mutations in the Apo B and Apo E genes that cause hypobetalipoproteinemia and dysbetalipoproteinemia, respectively. Mutations that are located within 10-15 nucleotides of the exon/intron boundary must be identified so that the homology region of the CMV continues with the sequence of the intron and not the exon. The genomic sequence of Apo E4 indicating the exon and intron boundaries is given in Paik et al., 1985, Proc. Natl. Acad. Sci. **82**, 3445, which is hereby incorporated by reference. The exon/intron boundaries of the Apo B gene are given in Table II along with the GENBANK accession numbers for the genomic sequence of Apo B.

6.2.4 Specific Alterations of the Apo A1 Gene

Apo A1 is a 243 amino acid protein. Amino acids 99-230 are encoded by six tandem duplications of a 66 base pair prototype sequence. The duplications are between 80 % and 64 % homologous to the consensus sequence. Without limitation as to theory the conformation of amino acids 120 to 230 is believed to be helical. The sequence of the amino acids is such that the helix is amphipathic, i.e., the helix has a hydrophobic face and a hydrophilic face, which contains polar amino acids.

The mutations most suitable for the practice of the invention are substitutions of polar amino acids by cysteine. Particularly suitable mutations are at arginine residues that are located next to other polar amino acids because the arginine codon used in apo A1 (CGC) can be converted into a TGC cysteine codon by a single base change. Thus, particularly suitable sites for mutations according to the present invention are arginine 149, 151, 153, 171 and 173.


6.3 FORMULATIONS SUITABLE FOR *IN VIVO* USE

The prior art formulations of CMV and a macromolecular carrier are of limited utility for *in vivo* use because of their low capacity for CMV and because the CMV is not protected from extracellular enzymes. The invention provides three alternative macromolecular carriers that overcome the limitations of the prior art. The carriers are polyethylenimine (PEI), aqueous-cored lipid vesicles, which are also termed unilamellar liposomes and lipid nanospheres.

Each of the carriers can be further provided with a ligand that is complementary to a cell-surface protein of the target cell. Such ligands are useful to increase both the amount and specificity of the uptake of CMV into the targeted cell. In one embodiment of the invention the target cell is a hepatocyte and the ligand is a galactose saccharide or lactose disaccharide that binds to the asialoglycoprotein receptor.

6.3.1 Polycationic Carriers

The invention can be practiced using any polycation that is non-toxic when administered to cells *in vitro* or to subjects *in vivo*. Suitable examples include polybasic amino acids such as polylysine, polyarginine, basic proteins such as histone H1, and synthetic polymers such as the branched-chain polyethylenimine:



The invention can be practiced with any branched chain polyethylenimine (PEI) having an average molecular weight of greater than about 500 daltons, preferably greater than between about 10 Kd and more preferably about 25 Kd (mass average molecular weight determined by light scattering). The upper limit of suitability is determined by the toxicity and solubility of the PEI. Toxicity and insolubility of molecular weights greater than about 1.3 Md makes such PEI material less suitable. The use of high molecular weight PEI as a carrier to transfect a cell with DNA is described in Boussif, O. et al., 1995, Proc. Natl. Acad. Sci. 92, 7297, which is hereby incorporated by reference. PEI solutions can be prepared according to the procedure of Boussif et al.

The CMV carrier complex is formed by mixing an aqueous solution of CMV and a neutral aqueous solution of PEI at a ratio of between 9 and 4 PEI nitrogens per CMV phosphate. In a preferred embodiment the ratio is 6. The complex can be formed, for

example, by mixing a 10 mM solution of PEI, at pH 7.0 in 0.15 M NaCl with CMV to form a final CMV concentration of between 100 and 500 nM.

In addition a ligand for a clathrin-coated pit receptor can be attached to the polycation or to a fraction of the polycations. In one embodiment the ligand is a saccharide or disaccharide that binds to the asialoglycoprotein receptor, such as lactose, galactose, or N-acetylgalactosamine. Any technique can be used to attach the ligands. The optimal ratio of ligand to polyethylene subunit can be determined by fluorescently labeling the CMV and injecting fluorescent CMV/molecular carrier/ligand complexes directly into the tissue of interest and determining the extent of fluorescent uptake according to the method of Kren et al., 1997, *Hepatology* **25**, 1462-1468.

Good results can be obtained using a 1:1 mixture of lactosylated PEI having a ratio of 0.4-0.8 lactosyl moieties per nitrogen and unmodified PEI. The mixture is used in a ratio of between 4 and 9 PEI nitrogens per CMV phosphate. A preferred ratio of oligonucleotide phosphate to nitrogen is 1:6. Good results can be obtained with PEIs having a mass average molecular weight of 25 Kd and 800 Kd which are commercially available from Aldrich Chemical Co., Catalog No. 40,872-7 and 18,197-8, respectively. Linear PEI such as that described in Ferrarri, S., et al., 1997, *Gene Therapy* **4**, 1100-1106 and sold under the trademark EXGEN 500™ is particularly suitable for the practice of the invention because of its lower toxicity compared to branched-chain PEI.

In an alternative embodiment the polycationic carrier can be a basic protein such as histone H1, which can be substituted with a ligand for a clathrin-coated pit receptor. A 1:1 (w/w) mixture of histone and CMV can be used to practice the invention.

6.3.2 Lipids that Are Useful in Carriers

The selection of lipids for incorporation into the lipid vesicle and lipid nanosphere carriers of the invention is not critical. Lipid nanospheres can be constructed using semi-purified lipid biological preparations, e.g., soybean oil (Sigma Chem. Co.) and egg phosphatidyl choline (EPC) (Avanti Polar Lipids). Other lipids that are useful in the preparation of lipid nanospheres and/or lipid vesicles include neutral lipids, e.g., dioleoyl phosphatidylcholine (DOPC), and dioleoyl phosphatidyl ethanolamine (DOPE), anionic lipids, e.g., dioleoyl phosphatidyl serine (DOPS) and cationic lipids, e.g., dioleoyl

trimethyl ammonium propane (DOTAP), dioctadecyldiamidoglycyl spermine (DOGS), dioleoyl trimethyl ammonium (DOTMA) and DOSPER (1,3-di-oleoyloxy-2-(6-carboxy-spermyl)-propyl-amide tetraacetate, commercially available from Boehringer-Mannheim). Additional examples of lipids that can be used in the invention can be found in Gao, X. and Huany, L., 1995, *Gene Therapy* **2**, 710. Saccharide ligands can be added in the form of saccharide cerebroside, e.g., lactosylcerebroside or galactocerebroside (Avanti Polar Lipids).

The particular choice of lipid is not critical. Hydrogenated EPC or lysolecithin can be used in place of EPC. DPPC (dipalmitoyl phosphatidylcholine), can be incorporated to improve the efficacy and/or stability of the delivery system.

6.3.3 The Construction of Lipid Nanosphere Carriers

Lipid nanospheres can be constructed by the following process. A methanol or chloroform methanol solution of phospholipids is added to a small test tube and the solvent removed by a nitrogen stream to leave a lipid film. A lipophilic salt of CMV is formed by mixing an aqueous saline solution of CMV with an ethanolic solution of a cationic lipid. Good results can be obtained when the cationic species are in about a 4 fold molar excess relative to the CMV anions (phosphates). The lipophilic CMV salt solution is added to the lipid film, vortexed gently followed by the addition of an amount of neutral lipid equal in weight to the phospholipids. The concentration of CMV can be up to about 3% (w/w) of the total amount of lipid.

After addition of the neutral lipid, the emulsion is sonicated at 4°C for about 1 hour until the formation of a milky suspension with no obvious signs of separation. The suspension is extruded through polycarbonate filters until a final diameter of about 50 nm is achieved. When the target cell is a reticuloendothelial cell the preferred diameter of the lipid nanospheres is about 100-200 nm. The CMV-carrying lipid nanospheres can then be washed and placed into a pharmaceutically acceptable carrier or tissue culture medium. The capacity of lipid nanospheres is about 2.5 mg CMV/ 500 μ l of a nanosphere suspension.

6.3.4 The Construction of Lipid Vesicles

A lipid film is formed by placing a chloroform methanol solution of lipid in a tube and removing the solvent by a nitrogen stream. An aqueous saline solution of CMV is added such that the amount of CMV is between 20% and 50% (w/w) of the amount of lipid, and the amount of aqueous solvent is about 80% (w/w) of the amount of lipid in the final mixture. After gentle vortexing the liposome-containing liquid is forced through successively finer polycarbonate filter membranes until a final diameter of about 50 nm is achieved. The passage through the successively finer polycarbonate filter results in the conversion of polylamellar liposomes into unilamellar liposomes, i.e., vesicles. When the target cell is a reticuloendothelial cell the preferred diameter of the lipid nanospheres is about 100-200 nm. The CMV-carrying lipid nanospheres can then be washed and placed into a pharmaceutically acceptable carrier or tissue culture medium.

The CMV are entrapped in the aqueous core of the vesicles. About 50% of the added CMV is entrapped.

A variation of the basic procedure comprises the formation of an aqueous solution containing a PEI/CMV condensate at a ratio of about 4 PEI imines per CMV phosphate. The condensate can be particularly useful when the liposomes are positively charged, i.e., the lipid vesicle contains a concentration of cations of cationic lipids such as DOTAP, DOTMA or DOSPER, greater than the concentration of anions of anionic lipids such as DOPS. The capacity of lipid vesicles is about 150 μ g CMV per 500 μ l of a lipid vesicle suspension.

In a preferred embodiment the lipid vesicles contain a mixture of the anionic phospholipid, DOPS, and a neutral lipid such as DOPE or DOPC. Other negatively charged phospholipids that can be used to make lipid vesicles include dioleoyl phosphatidic acid (DOPA) and dioleoyl phosphatidyl glycerol (DOPG). In a more preferred embodiment the neutral lipid is DOPC and the ratio of DOPS:DOPC is between 2:1 and 1:2 and is preferably about 1:1. The ratio of negatively charged to neutral lipid should be greater than 1:9 because the presence of less than 10% charged lipid results in instability of the lipid vesicles because of vesicle fusion.

A particular lipid vesicle formulation can be tested by using the formulation to transfect a target cell population with a plasmid of about 5.0 kb in length that expresses

some readily detectable product in the transfected target cell. Lipid vesicles can be used to transfect a cell with the plasmid if the plasmid is condensed with PEI at an imine:phosphate ratio of about 9-4:1. The capacity of the lipid vesicle formulation to transfect a cell with a plasmid is indicative of the formulation's capacity to introduce a CMV into a cell and effect a transmutation.

Certain lipids, particularly the polycationic lipids, can be toxic to certain cell lines and primary cell cultures. The formulation of the lipid vesicles should be adjusted to avoid such toxic lipids.

Ligands for clathrin-coated pit receptors can be introduced into the lipid vesicles by a variety of means. Cerebrosides, such as lactocerebroside or galactocerebroside can be introduced into the lipid mixture and are incorporated into the vesicle to produce a ligand for the asialoglycoprotein receptor.

In an alternative embodiment the lipid vesicle further comprises an integral membrane protein that inserts itself into the lipid bilayer of the vesicle. In a specific embodiment the protein is a fusogenic (F-protein) from the virus alternatively termed Sendai Virus or Hemagglutinating Virus of Japan (HVJ). The preparation and use of F-protein containing lipid vesicles to introduce DNA into liver, myocardial and endothelial cells have been reported. See, e.g., U.S. Patent No. 5,683,866, International Application PCT JP97/00612 (published as WO 97/31656). See also, Ramani, K., et al., 1996, FEBS Letters **404**, 164-168; Kaneda, Y., et al., 1989, J. Biol. Chem. **264**, 121126-12129; Kaneda, Y., et al., 1989, Science **243**, 375; Dzau, V.J., et al., Proc. Natl. Acad. Sci. **93**, 11421-11425; Aoki, M., et al., 1997, J.Mol.Cardiol. **29**, 949-959.

6.4 THE USE OF THE FORMULATIONS *IN VIVO*

The CMV of the invention can be parenterally administered directly to the target organ at a dose of between 50 and 250 $\mu\text{g/gm}$. When the target organ is the liver muscle or kidney, the CMV/macromolecular carrier complex can be injected directly into the organ. When the target organ is the liver, intravenous injection into the hepatic or portal veins of a liver, having temporarily obstructed circulation can be used. Alternatively the CMV/macromolecular complex can further comprise a hepatic targeting ligand, such as a

lactosyl or galactosyl saccharide, which allows for administration of the CMV/macromolecular complex intravenously into the general circulation.

When the target organ is the lung or a tissue thereof, e.g., the bronchiolar epithelium CMV/macromolecular complex can be administered by aerosol. Small particle aerosol delivery of liposomal/DNA complexes is described in Schwarz L.A., et al., 1996, Human Gene Therapy 7, 731-741.

When the target organ is the vascular endothelium, as for example in von Willebrand's Disease, the CMV/macromolecular complex can be delivered directly into the systemic circulation. Other organs can be targeted by use of liposomes that are provided with ligands that enable the liposome to be extravasated through the endothelial cells of the circulatory system.

For enzymatic defects, therapeutic effects can be obtained by correcting the genes of about 1% of the cells of the affected tissue. In a tissue in which the parenchymal cells have an extended life, such as the liver, treatments with CMV can be repeatedly performed to obtain an increased therapeutic effect.

7. EXAMPLES

7.1 CMV/MACROMOLECULAR CARRIER COMPLEXES

7.1.1 Lipid Nanospheres

Materials

Egg phosphatidylcholine (EPC), DOTAP and galactocerebroside (Gc) (Avanti Polar Lipids); soybean oil (Sigma Chemical Co.); dioctadecyldiamidoglycyl spermine (DOGS®) (Promega).

Methods

EPC, DOTAP and Gc were previously dissolved at defined concentrations in chloroform or anhydrous methanol and stored in small glass vials in desiccated containers at -20°C until use. EPC (40-45 mg), DOTAP (200 µg) and Gc (43 µg) solutions were aliquoted into a small 10 x 75 mm borosilicate tube and solvents removed under a stream of nitrogen. CMV were diluted in 0.15 M NaCl (~80-125 µg/250-300 µl); DOGS (as a 10 mg/ml solution in ethanol) was diluted into 250-300 µl 0.15 M NaCl at 3-5 times the weight of added CMV. The two solutions were mildly vortexed to mix contents and then CMV solution was added slowly to the DOGS solution. The contents were mixed by gentle tapping and inverting the tube a few times. The DOGS-complex solution was added to the dried lipids followed by soybean oil (40-45 mg), the mixture was vortexed on high for a few seconds and bath sonicated in a FS-15 (Fisher Scientific) bath sonicator for ~1 hr in a 4°C temperature controlled room. Occasionally, the tube was removed from the bath and vortexed. When a uniform looking, milky suspension was formed (with no obvious separation of oil droplets), it was extruded through a series of polycarbonate membranes down to a pore size of 50 nm. Preparations were stored at 4°C until use and vortexed before use.

7.1.2 Negatively Charged Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl phosphatidylserine (DOPS), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids).

Methods

DOPS, DOPC and Gc at a molar ratio of 1:1:0.16 (500 μg total lipid) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. The CMV were diluted in 500 μl of 0.15 M NaCl (approximately 100-250 μg /500 μl). The solution was added to the lipid film at room temperature. Lipids were dispersed entirely by alternate mild vortexing and warming (in a water bath at 37-42°C). After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μm) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation, lipid vesicles were stored at 4°C until use. Under these conditions the lipid vesicles were stable for at least one month. The final product can be lyophilized.

7.1.3 Neutral Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl phosphatidylethanolamine (DOPE), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids).

Methods

DOPC, DOPE and Gc (1:1:0.16 molar ratio) or DOPC:Gc (1:0.08) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. The oligonucleotides (or chimeric molecules) were diluted in 500 μl of 0.15 M NaCl (approximately 100-250 μg /500 μl). The solution was added to the lipid film at room temperature. Lipids were dispersed entirely by alternate mild vortexing and warming (in a water bath at 37-42°C). After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μm) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation, lipid vesicles were stored at 4°C until use. The size of the lipid vesicles of the preparation was stable for about 5 days.

7.1.4 Positively Charged Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl trimethylammonium propane (DOTAP), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids). Polyethylenimine (PEI) (M.W. 800 Kd), Fluka Chemicals.

Methods

DOPC, DOTAP and Gc (6:1:0.56 molar ratio) (500 μ g total lipid) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. PEI was diluted to a concentration of 45 mg/100 ml using water. pH of the solution was adjusted to ~7.6 using HCl. This PEI stock solution was prepared fresh each time and was equivalent to approximately 50 nmol amine/ μ l. CMV were diluted into 0.15 M NaCl at a concentration of ~125 μ g in 250 μ l. PEI was further diluted into 250 μ l 0.15 M NaCl so that approximately 4 moles of PEI amine were present per mole of oligonucleotide/chimeric phosphate. PEI solution was added drop-wise to the CMV solution (both at room temperature) and vortexed for 5-10 minutes. The PEI-complex solution was then added to the lipid film and the lipids dispersed as described above. After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μ m) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation lipid vesicles were stored at 4°C until use. Under these conditions the lipid vesicles were stable for at least one month. For longer and improved stability the final product can be lyophilized.

7.1.5 Lactosylated-PEI/PEI Complexes

PEI (25 kDa) was purchased from Aldrich Chemical (Milwaukee, WI). PEI (800 kDa) was purchased from Fluka chemicals (Ronkonkoma, NY, USA). Lactosylation of the PEI was carried out by modification of a previously described method for the conjugation of oligosaccharides to proteins. Briefly, 3 to 5 ml of PEI (0.1 to 1.2 M_{monomer}) in ammonium acetate (0.2 M) or Tris buffer (0.2 M) (pH 7.6) solution was incubated with 7 to 8 mg of sodium cyanoborohydride (Sigma Chemical Co., St. Louis, MO) and

approximately 30 mg of lactose monohydrate (Sigma Chemical Co., St. Louis, MO). Reaction was carried out in polypropylene tubes, tightly capped in a 37°C shaking water bath. After 10 days the reaction mixture was dialyzed against distilled water (500 ml) for 48 h with 1 to 2 changes of water. The purified complex was sterile filtered through 0.2 μ m filter and stored at 4°C. The amount of sugar (as galactose) associated with PEI was determined by the phenol-sulphuric acid method.

The number of moles of free amine (primary + secondary) in the lactosylated PEI was determined as follows: a standard curve was set up using a 0.02M stock solution of PEI; several aliquots of the stock were diluted to 1ml using deionized water in glass tubes, then 50 μ l of Ninhydrin reagent (Sigma Chemical Co., St. Louis, Mo) was added to each tube and vortexed vigorously for 10 sec. Color development was allowed to proceed at room temperature for 10 to 12 min. and then O.D. was read (within 4 minutes) at 485 nm on a Beckman DU-64 spectrophotometer. 20 to 50 μ l aliquots of the L-PEI samples were treated as above and the number of moles of free amine was determined from the standard curve. Lactosylated-PEI (L-PEI) complexes were prepared as follows: an equivalent of 3 mmol of amine as L-PEI and 3 mmol of amine as PEI, per mmol of RNA/DNA phosphate, were mixed together and diluted in 0.15M NaCl as required; the mixture was added dropwise to a solution of the chimeric and vortexed for 5 min.

To verify complete association of the chimeric oligonucleotides with PEI or L-PEI, gel analysis (4% LMP agarose) of the uncomplexed and complexed chimerics was performed. To determine the degree of protection against nuclease degradation provided by complexation of the chimerics, samples were treated with RNase and DNase. After a chloroform phenol extraction, the complexes were dissociated using heparin (50 units/ μ g nucleic acid) and the products analyzed on a 4% LMP agarose gel.

7.2 DEMONSTRATION OF PEI/CMV MEDIATED ALTERATION OF RAT AND HUMAN FACTOR IX

Materials. Fetal bovine serum was obtained from Atlanta Biologicals, Inc. (Atlanta, GA). The terminal transferase, fluorescein-12-dUTP, Expand™ high fidelity PCR system, dNTPs and high pure PCR template preparation kit were obtained from Boehringer Mannheim Corp. (Indianapolis, IN). Reflection™ NEF-496 autoradiography

film and Reflection™ NEF-491 intensifying screens were from DuPont NEN® Research Products (Boston, MA). Polyethylenimine (PEI) 800 kDa was obtained from Fluka Chemical Corp. (Ronkonkoma, NY). The [γ -³²P]ATP was obtained from ICN Biochemicals, Inc. (Costa Mesa, CA). pCR™2.1 was obtained from Invitrogen (San Diego, CA). OPTIMEM™, Dulbecco's modified Eagle's medium, William's E medium and oligonucleotides 365-A and 365-C were from Life Technologies, Inc. (Gaithersburg, MD). Spin filters of 30,000 mol wt cutoff were purchased from Millipore Corp. (Bedford, MA). Dil and SlowFade™ antifade mounting medium were obtained from Molecular Probes, Inc. (Eugene, OR). T4 polynucleotide kinase was purchased from New England Biolabs, Inc. (Beverly, MA). MSI MagnaGraph membrane was purchased from Micron Separations, Inc. (Westboro, MA). The primers used for PCR amplification were obtained from Oligos Etc., Inc. (Wilsonville, OR). Tetramethylammonium chloride was purchased from Sigma Chemical Company (St. Louis, MO). All other chemicals were molecular biology or reagent grade and purchased from Aldrich Chemical Company (Milwaukee, WI), Curtin Matheson Scientific, Inc. (Eden Prairie, MN), and Fisher Scientific (Itasca, IL).

Oligonucleotide synthesis. Chimeric RNA/DNA oligonucleotides HIXF, RIXF and RIXR were synthesized. The CMV were prepared with DNA and 2'-O-methyl RNA phosphoramidite nucleoside monomers on an ABI 394 synthesizer. The DNA phosphoramidite exocyclic amine groups were protected with benzoyl (adenosine and cytidine) and isobutyryl (guanosine). The protective groups on the 2'-O-methyl RNA phosphoramidites were phenoxyacetyl for adenosine, isobutyryl for cytidine, and dimethylformamide for guanosine. The base protecting groups were removed following synthesis by heating in ethanol/concentrated ammonium hydroxide for 20 h at 55°C. The crude oligonucleotides were electrophoresed on 15% polyacrylamide gels containing 7 M urea, and the DNA visualized using UV shadowing. The chimeric molecules were eluted from the gel slices, concentrated by precipitation and desalted using G-25 spin columns. Greater than 95% of the purified oligonucleotides were full length.

The sequence of the wild type and "mutant" rat Factor IX are

(SEQ ID No. 27) 365
wt AAA GAT TCA TGT GAA GGA GAT AGT GGG GGA CCC CAT GTT
 Lys Asp Ser Cys Glu Gly Asp Ser Gly Gly Pro His Val
 (SEQ ID No. 28)

(SEQ ID No. 29)
mt AAA GAT TCA TGT GAA GGA GAT CGT GGG GGA CCC CAT GTT
 Arg

The structure of the RIXR, RIXF and HIXR CMV is as follows:

Chimeric Oligonucleotides

RIXR (SEQ ID No. 30)

TGCGCG-ccccagggggTGCTAgaggaaguguT
 T T
 T T
 TCGCGC GGGGTCCCCCACGATCTCCTTCACAT
 3' 5'

RIXR_C (SEQ ID No. 31)

TGCGCG-acacuuccucTAGCAccccccuggggT
 T T
 T T
 TCGCGC TGTGAAGGAGATCGTGGGGGACCCCT
 3' 5'

RIXF (SEQ ID No. 32)

TGCGCG-acacuuccucTAGCAccccccuggggT
 T T
 T T
 TCGCGC TGTGAAGGAGATCGTGGGGGACCCCT
 3' 5'

HIXF (SEQ ID No. 33)
 TGC GCG-acaguuccucTAGCAccccccuggggT
 T T
 T T
 TCGCGC TGTCAAGGAGATCGTGGGGGACCCCT
 3' 5'

Uppercase letters are deoxyribonucleotides, lower case letters are 2'OMe-ribonucleotides. The nucleotide of the heterologous region is underlined.

Cell Culture, transfections and hepatocyte isolation. HuH-7 cells were maintained in Dulbecco's modified Eagle's medium containing 10% (vol/vol) heat inactivated fetal bovine serum in a humidified CO₂ atmosphere at 37°C. Twenty four hours prior to transfection 1 x 10⁵ cells were plated per 35 mm culture dish. At the time of transfection, the cells were rinsed twice with OPTIMEM™ media and transfections were performed in 1 ml of the same media. Eighteen hours after transfection, 2 ml of Dulbecco's modified Eagle's medium containing 20% (vol/vol) heat inactivated fetal bovine serum was added to each 35 mm dish and the cells maintained for an additional 30 h prior to harvesting for DNA isolation. A PEI (800 kDa) 10 mM stock solution, pH 7.0, was prepared. Briefly, the chimeric oligonucleotides were transfected with 10 mM PEI at 9 equivalents of PEI nitrogen per chimeric phosphate in 100 µl of 0.15 M NaCl at final concentrations of either 150 nM (4 µg), 300 nM (8 µg) and 450 nM (12 µg). After 18 h, an additional 2 ml of medium was added and reduced the chimeric concentrations to 50 nM, 100 nM, and 150 nM, respectively, for the remaining 30 h of culture. HuH-7 vehicle control transfections utilized the same amount of PEI as was used in the HuIXF transfections, but substituted an equal volume of 10 mM Tris-HCl pH 7.6 for the oligonucleotides.

Primary rat hepatocytes were isolated from 250 g male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc., Indianapolis, IN) by a two step collagenase perfusion as previously described (Fan et al., Oncogene 12:1909-1919, 1996, which is hereby incorporated by reference) and plated on Primaria™ plates at a density of 4 x 10⁵ cells per 35 mm dish. The cultures were maintained in William's E medium supplemented with 10% heat inactivated FBS, 26 mM sodium bicarbonate, 23 mM HEPES, 0.01 U/ml

insulin, 2 mM L-glutamine, 10 nM dexamethasone, 5.5 mM glucose, 100 U/ml penicillin and 100 U/ml streptomycin. Twenty four hours after plating, the hepatocytes were washed twice with the same medium and 1 ml of fresh medium added and the cells transfected using PEI/chimeric oligonucleotide complexes at the identical concentrations as for the HuH-7 cells. After 18 h, an additional 2 ml of the medium was added and the cells harvested 6 or 30 h later.

Direct injection of chimeric oligonucleotides into liver. Male Sprague-Dawley rats (~175 g) were maintained on a standard 12 h light-dark cycle and fed *ad libitum* standard laboratory chow. The rats were anesthetized, a midline incision made the liver exposed. A clamp was placed on the hepatic and portal veins as they enter the caudate lobe, and 75 μ g of the 1:9 chimeric/PEI complex was injected in a final volume of 250 - 300 μ l directly into the caudate lobe. The lobe remained ligated for 15 min and then blood flow was restored by removing the clamp. After suturing the incision the animals were allowed to recover from the anesthesia and given food and water *ad libitum*. Vehicle controls were done substituting an equal volume of Tris-HCl pH 7.6 for the chimeric oligonucleotides. Twenty-four and 48 h post-injection the animals were sacrificed, the caudate lobe removed and the tissue around the injection site dissected for DNA isolation. DNA was isolated and the terminal exon of the rat factor IX gene was amplified by PCR.

Nuclear uptake of the chimeric molecules. Chimeric duplexes were 3' end-labeled using terminal transferase and fluorescein-12-dUTP according to the manufacturer's recommendation, and were then mixed with unlabeled oligonucleotides at a 2:3 ratio. Transfections were performed as described above and after 24 h the cells were fixed in phosphate buffered saline, pH 7.4, containing 4% paraformaldehyde (wt/vol) for 10 min at room temperature. Following fixation, the cells were counterstained using a 5 μ M solution of Dil in 0.32 M sucrose for 10 min according to the manufacturer's recommendation. After washing with 0.32 M sucrose and then phosphate buffered saline, pH 7.4, the cells were coverslipped using SlowFade™ antifade mounting medium in phosphate buffered saline and examined using a MRC1000 confocal microscope (BioRad, Inc., Hercules, CA). The caudate lobes of liver *in situ* were injected with fluorescently-labeled chimerics as described above and harvested 24 h post-

injection. The lobes were bisected longitudinally, embedded using OCT and frozen. Cryosections were cut ~10 μ m thick, fixed for 10 min at room temperature using phosphate buffered saline, pH 7.4, containing 4% paraformaldehyde (wt/vol). Following fixation, the cells were counterstained using a 5 μ M solution of Dil in 0.32 M sucrose for 10 min according to the manufacturer's recommendation. After washing with 0.32 M sucrose and then phosphate buffered saline, pH 7.4, the sections were coverslipped using SlowFade™ antifade mounting medium and examined using a MRC1000 confocal microscope (BioRad, Inc.). The collection series for the fixed cells and sectioned tissue were made at 1 μ m steps to establish the presence of the chimeric in the nucleus.

DNA isolation and cloning. The cells were harvested by scrapping 24 and 48 h after transfection. Genomic DNA larger than 100-150 base pairs was isolated using the high pure PCR template preparation kit according to the manufacturer's recommendation. PCR amplification of a 317-nt fragment of the eighth exon in the human liver factor IX gene was performed with 500 ng of the isolated DNA. The primers used were 5'-CATTGCTGACAAGGAATACACGAAC-3' (SEQ ID No. 34) and 5'-ATTGCCTTTCATTGCACACTCTTC-3' (SEQ ID No. 35) corresponding to nucleotides 1008-1032 and 1300-1324, respectively, of the human factor IX cDNA. Primers were annealed at 58°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelity™ polymerase. PCR amplification of a 374-nt fragment of the rat factor IX gene was performed with 500 ng of the isolated DNA from either the primary hepatocytes or liver caudate lobe. The primers used were 5'-ATTGCCTTGCTGGAACTGGATAAC-3' (SEQ ID No. 36) and 5'-TTGCCTTTCATTGCACATTCTTCAC-3' (SEQ ID No. 37) corresponding to nucleotides 433-457 and 782-806, respectively, of the rat factor IX cDNA. Primers were annealed at 59°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelity™ polymerase. The PCR amplification products from both the human and rat factor IX genes were subcloned into the TA cloning vector pCR™2.1 according to the manufacturer's recommendations, and the ligated material used to transform frozen competent *Escherichia coli*.

Colony hybridization and sequencing. Eighteen to 20 h after plating, the colonies were lifted onto MSI MagnaGraph nylon filters, replicated and processed for hybridization according to the manufacturer's recommendation. The filters were hybridized for 24 h with 17 mer oligonucleotide probes 365-A (5'-AAGGAGATAGTGGGGGA-3') (SEQ ID No. 38) or 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 39), where the underlined nucleotide is the target of the mutagenesis. The probes were ³²P-end-labeled using [γ -³²]ATP (>7,000 Ci/mmol) and T4 polynucleotide kinase according to the manufacturer's recommendations. Hybridizations were performed at 37°C in 2X sodium chloride sodium citrate containing 1% SDS, 5X Denhardt's and 200 μ g/ml denatured sonicated fish sperm DNA. After hybridization, the filters were rinsed in 1X sodium chloride phosphate EDTA, 0.5% SDS and then washed at 54°C for 1 h in 50 mM Tris-HCl, pH 8.0 containing 3 M tetramethylammonium chloride, 2 mM EDTA, pH 8.0, 0.1% SDS. Autoradiography was performed with NEN® Reflection film at -70°C using an intensifying screen. Plasmid DNA was prepared from colonies identified as hybridizing with 365-A or 365-C using Qiagen minprep kit (Chatsworth, CA) and subjected to automatic sequencing using the mp13 reverse primer on an ABI 370A sequencer (Perkin-Elmer, Corp., Foster City, CA).

Results *In Vivo*

Chimeric oligonucleotides were fluorescein-labeled and used to determine whether direct injection into the caudate lobe of the liver was feasible. The results indicated that the hepatocytes adjacent to the injection site within the caudate lobe showed uptake of the fluorescently-labeled chimeric molecules similar to that observed in isolated primary hepatocytes and HuH-7 cells. Although some punctate material was present in the cytoplasm, the labeled material was detected primarily in the nucleus. In fact, only nuclear labeling was observed in hepatocytes farthest from the injection site. The unlabeled PEI/RIXF chimeric complexes and vehicle controls were injected directly into the caudate lobe using the same protocol and the animals sacrificed 24 and 48 h post-injection. Liver DNA was isolated as described in Methods, subjected to PCR amplification of a 374 nt sequence spanning the targeted nt exchange site. Following

subcloning and transformation of *Escherichia coli* with the PCR amplified material, duplicate filter lifts of the transformed colonies were performed. The filters were hybridized with ³²-labeled 17-mer oligonucleotides specific for either 365-A (wild-type) or 365-C (factor IX mutation) and processed post-hybridization as described in Methods. Rats which received direct hepatic injection of the RIXF chimeric molecules exhibited a A→C conversion frequency of ~1% at both 24 and 48 h. In contrast, the vehicle controls showed no hybridization with the 365-C probe. Colonies that hybridized with the 365-C probe from the RIXF treated animals were cultured, the plasmid DNA isolated and subjected to sequencing to confirm the A→C conversion. The ends of the amplified 374-nt fragment correspond exactly with the primers and the only nucleotide change observed was an A→C at the targeted exchange site.

7.3 DEMONSTRATION OF LACTOSYLATED-PEI/CMV MEDIATED ALTERATION OF RAT FACTOR IX

7.3.1 Results

CMV complexed to a mixture of lactosylated-PEI and PEI was prepared using the RIXR oligonucleotide as described in Section 6.1.5 above. A CMV directed to the complementary strand of the same region of the factor IX was also constructed (RIXR_C).

Conversion of the targeted nucleotide at Ser³⁶⁵ by the chimeric oligonucleotides

The nuclear localization of the fluorescently-labeled chimeric molecules indicated efficient transfection in the isolated rat hepatocytes. The cultured hepatocytes were then transfected with the unlabeled chimeric molecules factor RIXR_C and RIXR at comparable concentrations using 800 kDa PEI as the carrier. Additionally, vehicle control transfections were performed simultaneously. Forty eight hours after transfection, the cells were harvested and the DNA isolated and processed for hybridization as described in Section 6.1.5. The A→C targeted nucleotide conversion at Ser³⁶⁵ was determined by hybridization of duplicate colony lifts of the PCR-amplified and cloned 374-nt stretch of exon 8 of the factor IX gene (Sarkar, B., Koeberl, D. D. & Somer, S. S., "Direct Sequencing of the activation peptide and the catalytic domain of the factor IX gene in six species," *Genomics*, 6, 133-143, 1990.) The 17 mer oligonucleotide probes used to distinguish between the wild-type 365-A (5'-AAGGAGATAGTGGGGGA-3') (SEQ ID No.

40) or converted 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 41) corresponded to nucleotides 710 through 726 of the cDNA sequence.

The overall frequency of conversion of the targeted nucleotide was calculated by dividing the number of clones hybridizing with the 365-C oligonucleotide by the total number of clones hybridizing with both oligonucleotide probes. The results are summarized in Table III for RIXR_C. A→C conversion at Ser³⁶⁵ was observed only in primary hepatocytes transfected with the RIXR or RIXR_C. Similar conversion frequencies were observed in hepatocytes transfected with RIXR or RIXR_C. Neither vehicle transfected cells nor those transfected with other chimeric oligonucleotides yielded any clones hybridizing with the 365-C oligonucleotide probe (unpublished observations). Additionally, no hybridization of the 365-C oligonucleotide probe was observed to clones derived from DNA isolated from untreated hepatocytes and PCR-amplified in the presence of 0.5 to 1.5 μg of the oligonucleotides. The A→C conversion rate in the isolated hepatocytes was also dose dependent using lactosylated PEI derivatives as described in Section 6.1.5 and was as high as 19%. RT-PCR and hybridization analysis of RNA isolated from cultured cells transfected in parallel with lactosylated PEIs demonstrated A→C conversion frequencies ranging from 11.9 to 22.3%.

Site-directed nucleotide exchange by chimeric oligonucleotides in intact liver

The fluorescein-labeled oligonucleotides were also used to determine cellular uptake of the chimeric molecules after direct injection into the caudate lobe of the liver. The results indicated that hepatocytes adjacent to the injection site in the caudate lobe showed uptake of the fluorescently-labeled chimerics similar to that observed in the isolated rat hepatocytes. Although some punctate material was present in the cytoplasm of the hepatocytes, the labeled material was primarily present in the nucleus. In fact, only nuclear labeling was observed in those areas farthest from the injection site. The unlabeled RIXR chimeric oligonucleotides and vehicle controls were then administered *in vivo* by tail vein injection of the 25 kDa PEI and liver tissue harvested 5 days post-injection. Liver DNA was isolated and subjected to PCR amplification of a 374-nt sequence spanning the targeted nucleotide exchange site, using the same primers as those used with the primary hepatocytes. Following subcloning and transformation of *E. coli*

with the PCR-amplified material, duplicate filter lifts of the transformed colonies were done. The filters were hybridized with the same ^{32}P -labeled 17-mer oligonucleotides specific for either 365-A (wild-type) or 365-C (mutant) and processed post-hybridization. Rats treated with 100 μg of the RIXR chimeric oligonucleotides exhibited an A \rightarrow C conversion frequency ranging from 13.9% to 18.9%, while those that received a total of 350 μg in two injections showed 40% conversion. In contrast, the vehicle controls showed no hybridization with the 365-C probe. RT-PCR hybridization of isolated RNA indicated A \rightarrow C conversion frequencies of 26.4% to 28.4% in the high dose livers. The APTT for vehicle-treated rats ranged from 89.7% to 181.9% of control values ($131.84\% \pm 32.89\%$), while the APTT for the oligonucleotide-treated animals ranged from 48.9% to 61.7% ($53.8\% \pm 4.8\%$).

The APTT times for a 1/10 dilution of rat test plasma in Hepes buffer (50 mM Hepes/100 mM NaCl/0.02% NaN_3 , pH 7.4) were determined for both normal ($n = 9$) and the double injected animals ($n = 3$). The factor IX activity of duplicate samples was determined from a log-log standard curve that was constructed from the APTT results for dilution (1:10 to 1:80) of pooled plasma from 12 normal male rats, 6-8 weeks old. The APTT results for the normal rats ranged from 89.7% to 181.9% of the control values (mean = $131.84\% \pm 32.89\%$), while the APTT results for the double injected animals ranged from 49.0% to 61.7% (mean $53.8\% \pm 5.8\%$). The APTT clotting time in seconds for the normal rats ranged from 60.9 seconds to 81.6 seconds (mean = 71.3 ± 7.3 seconds) while the APTT times ranged from 92.3 seconds to 98.6 seconds (mean = 96.3 ± 2.9 seconds) for the double-infected rats.

Sequence analysis of the mutated factor IX gene in isolated hepatocytes and intact liver

Direct sequencing of the wild-type and mutated genes was performed to confirm the results from the filter hybridizations in both the *in vitro* and *in vivo* studies. At least 10 independent clones hybridizing to either 365-A or 365-C from the intact liver or isolated hepatocytes were analyzed. The results of the sequencing indicated that colonies hybridizing to 365-A (Fig. 6, top panel) exhibited the wild-type IX sequence, i.e. an A at Ser³⁶⁵ of the reported cDNA sequence. In contrast, those colonies derived from the factor RIXR_C transfected primary hepatocytes hybridizing to the 365-C oligonucleotide probe

converted to a C at Ser³⁶⁵. The same A→C conversion at Ser³⁶⁵ was observed in the clones derived from the transfected rat liver that hybridized with the 17 mer 365-C oligonucleotide probe. The entire 374-nt PCR amplified region of the factor IX gene was sequenced for all the clones and no alteration other than the indicated changes at Ser³⁶⁵ was detected. Finally, the start and end points of the 374-nt PCR amplified genomic DNA derived from both the primary hepatocytes and the intact liver corresponded exactly to those of the primers used for the amplification process, indicating that the cloned and sequenced DNA was derived from genomic DNA rather than nondegraded chimeric oligonucleotides.

Table III Percent A-C conversion at Ser³⁶⁵ of rat factor IX genomic DNA by colony lift hybridizations

PEI Deliver System		365-C clones	Total clones	A-C (%)
PEI 800 kDa ¹	<u>Concentration</u>			
<i>In vitro</i>	150 nM	24	572	4.2
	300	31	367	8.5
	450	63	502	12.5
Lac-PEI 800 kDa				
<i>In vitro</i>	90	18	337	5.3
	180	34	300	11.3
	270	47	253	18.6
Lac-PEI 25 kDa				
<i>In vitro</i>	90	28	527	5.3
	180	53	417	12.7
	270	60	305	19.7
Lac-PEI 25 kDa ²	<u>Dose</u>			
<i>In vivo</i> x1	100 µg	24	166	14.5
		71	386	18.4
		50	360	13.9
Lac-PEI 25 kDa				
<i>In vivo</i> x2	350 µg	237	601	39.4
		228	563	40.5
		271	678	40.0

¹The data shown for the primary hepatocyte transfections represents a mean of two experiments.

²The *in vivo* chimeric/PEI complexes were administered in a volume of 300 µl of 5% dextrose by tail vein injection. The results of three animals at each dose are shown individually.

7.3.2 Materials and Methods

In vivo delivery of the chimeric oligonucleotides. Male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc.) (~50 g) were maintained on a standard 12 h light-dark cycle and fed *ad libitum* standard laboratory chow. Vehicle controls and lactosylated 25 kDa PEI at a ratio of 6 equivalents of PEI nitrogen per chimeric phosphate were administered in 300 μ l of 5% dextrose (Abdallah, B. et al., "A powerful nonviral vector for *in vivo* gene transfer into the adult mammalian brain: polyethylenimine; *Human Gene Therapy*, 7, 1947-1954, 1996.). The aliquots were administered by tail vein injection either as a single dose of 100 μ g or divided dose of 150 μ g and 200 μ g on consecutive days. Five days post-injection, liver tissue was removed for DNA and RNA isolation. DNA was isolated as previously described (Kren, B. T., Trembley, J. H. & Steer, C. J., "Alterations in mRNA stability during rat liver regeneration," *Am. J. Physiol.*, 270, G763-G777, 1996) for PCR amplification of exon 8 of the rat factor IX gene. RNA was isolated for RT-PCR amplification of the same region as the genomic DNA using RNAexol and RNAmate (Intermountain Scientific Corp., Kaysville, UT) according to the manufacturer's protocol.

Factor IX activity assay. Blood samples from vehicle (n = 9) and oligonucleotide-treated (n = 3) rats were collected 20 days after the second tail vein injection in 0.1 vol. of 0.105 M sodium citrate/citric acid. After centrifugation at 2,500 x g and then 15,000 x g the resulting plasma was stored at -70°C. The factor IX activity was determined from activated partial thromboplastin time (APTT) assays. Briefly, 50 μ l of APTT reagent (DADE, Miami, FL), 50 μ l of human factor IX-deficient plasma (George King Biomedical, Overland, KS), and 50 μ l of 1/10 dilution of rat test plasma in Hepes buffer (50 mM Hepes/100 mM NaCl/0.02% NaN₃, pH 7.4) were incubated at 37°C for 3 min in an ST4 coagulometer (American Bioproducts, Parsippany, NJ). Clotting was initiated by addition of 50 μ l of 33 mM CaCl₂ in Hepes buffer. Factor IX activity of duplicate samples was determined from a log-log standard curve constructed from the APTT results for dilution (1:10 to 1:80) of pooled plasma from normal male rats (n = 12).

DNA/RNA isolation and cloning. The cells were harvested by scrapping 48 h after transfection. Genomic DNA larger than 100-150 base pairs was isolated using the high

pure PCR template preparation kit (Boehringer Mannheim, Corp., Indianapolis, IN). RNA was isolated using RNAzol™ B (Tel-Test, Inc., Friendswood, TX), according to the manufacturer's protocol. PCR amplification of a 374-nt fragment of the rat factor IX gene was performed with 300 ng of the isolated DNA from either the primary hepatocytes or liver tissue. The primers were designed as 5'-ATTGCCTTGCTGGAAGTGGATAAAC-3' (SEQ ID No. 42) and 5'TTGCCTTTCATTGCACATTCTTCAC-3' (SEQ ID No. 43) (Oligos Etc., Wilsonville, OR) corresponding to nucleotides 433-457 and 782-806, respectively, of the rat factor IX cDNA. Primers were annealed at 59°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelity™ polymerase (Boehringer Mannheim, Corp.). The PCR amplification products from both the hepatocytes and intact liver factor IX genes were subcloned into the TA cloning vector pCR™2.1 (Invitrogen, San Diego, CA), and the ligated material used to transform frozen competent *E. coli*. To rule out PCR artifacts 300 ng of control DNA was incubated with 0.5, 1.0 and 1.5 µg of the oligonucleotide prior to the PCR-amplification reaction. Additionally, 1.0 µg of the chimeric alone was used as the "template" for the PCR amplification.

RT-PCR amplification was done utilizing the Titian™ one tube RT-PCR system (Boehringer Mannheim, Corp.) According to the manufacturer's protocol using the same primers as those used for the DNA PCR amplification. To rule out DNA contamination, the RNA samples were treated with RQ1 DNase free RNase (Promega Corp., Madison, WI) and RT-PCR negative controls of RNased RNA samples were performed in parallel with the RT-PCR reaction. Each of the PCR reactions were ligated into the same TA cloning vector and transformed into frozen competent *E. coli*.

Colony hybridization and sequencing. Eighteen to 20 h after plating, the colonies were lifted onto MSI MagnaGraph nylon filters, replicated and processed for hybridization according to the manufacturer's recommendation. The filters were hybridized for 24 h with 17 mer oligonucleotide probes 365-A (5'AAGGAGATAGTGGGGGA-3') (SEQ ID No. 44) OR 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 45) (Life technologies, Inc., Gaithersburg, MD), where the underlined nucleotide is the target for

mutagenesis. The probes were ^{32}P -end-labeled using (γ - ^{32}P) ATP ($> 7,000$ Ci/mmol) and T4 polynucleotide kinase (New England Biolabs, Inc., Beverly MA). Hybridizations were performed at 37°C in 2X sodium chloride sodium citrate containing 1% SDS, 5X Denhardt's and 200 $\mu\text{g/ml}$ denatured sonicated fish sperm DNA. After hybridization, the filters were rinsed in 1X sodium chloride sodium phosphate EDTA, 0.5% SDS and then washed at 54°C for 1 h in 50 mM Tris-HCl, pH 8.0 containing 3 M tetramethylammonium chloride, 2 mM EDTA, pH 8.0, 0.1% SDS (Melchior, W. B. & Von Hippel, P. H. "Alteration of the relative stability of dA.dT and dG.dC base pairs in DNA," Proc. Natl. Acad. Sci. USA, **70**, 298-302, 1973.). Autoradiography was performed with NEN[®]Reflection film at -70°C using an intensifying screen. Plasmid DNA was prepared from colonies identified as hybridizing with 365-A or 365-C using Qiagen miniprep kit (Chatsworth, CA) and subjected to automatic sequencing using the mp13 forward and reverse primers as well as a gene specific primer, 5'GTTGACCGAGCCACATGCCTTAG-3' (SEQ ID No. 46) corresponding to nucleotides 616 to 638 of the rat factor IX cDNA using an ABI 370A sequencer (Perkin-Elmer, Corp., Foster City, CA).

7.4 EXAMPLES OF CMV USEFUL FOR THE REDUCTION OF LDL LEVELS IN HUMANS

A CMV suitable for the modification of Apo B having a sequence comprising the sequence of SEQ ID No: 5 is given below.

Apo B 41/UR (mut \rightarrow WT)

(SEQ ID No. 47)

```

u GCGCG gac ccg acc gaa uuc ggu aac ugu au
u                                     u
u                                     u
u CGCGC CTG GGC TGG CTT AAG CCA TTG ACA Tu
      3' 5'

```

A CMV suitable for the modification of Apo B having a sequence comprising the sequence of SEQ ID No: 12 is given below.

Apo B 5/U88 (mut→WT)

(SEQ ID No. 48)

```

u GCGCG cug uuc aaa gug uaC GGA TCC ucu uug acu gac gau
u                                     u
u                                     u
u CGCGC GAC AAG TTT CAC ATG CCT AGG AGA AAC TGA CTG CTu
    3' 5'

```

7.5 CORRECTION OF A CRIGLER-NAJJAR-LIKE MUTATION IN THE GUNN RAT

Mutant rats with hyperbilirubinemia, termed Gunn rats, have a single nucleotide deletion in the gene encoding bilirubin-uridinediphosphoglucuronate glucuronosyltransferase (*UGT1A1*). Roy Chowdhury, J., et al., 1991, J. Biol. Chem. **266**, 18294. Human patients with Crigler-Najjar syndrome type I also have mutations of the *UGT1A1* gene, resulting in life-long hyperbilirubinemia and consequent brain damage. Bosma, P.J., et al., 1992, FASEB J. **6**, 2859; Jansen, P.L.M., et al., Progress In Liver Diseases, **XIII**, Boyer, J.L., & Ockner, R.K., editors (W.B. Saunders, Phil. 1995), pp 125-150. The structure of CN3, a CMV designed to correct the Gunn rat mutation is given below.

CN3 (mut→WT)

(SEQ ID No. 49)

```

T GCGCG gg gac uua caG GAC CTT TAC uga ctt cua T
T                                     T
T                                     T
T CGCGC CC CTG AAT GTC CTG GAA ATG ACT GCC GAT T
    3' 5'

```

Gunn rat primary cultured hepatocytes were treated with 150 nM CN3 according to the above protocol except that the carrier was either the negatively charged glycosylated lipid vesicles of section 6.2.2 or a lactosylated-PEI carrier at a ratio of

oligonucleotide phosphate to imine of 1:4 . The results were 8.5% conversion with the negatively charged liposome and 3.6% conversion with lactosylated-PEI carrier.

Gunn rats were injected with 1 mg/Kg of CN3 complexed with either 25 kDa Lac-PEI or complexed with negatively charged Gc lipid vesicles (Gc-NLV) as described above. The rate of gene conversion was determined by cloning and hybridization according to the procedure described for factor IX. The results shown below indicate that between about 15% and 25% of the copies of the *UGT1A1* gene were converted.

**Frequency of Insertion of G at nucleotide 1239 of the UGT-1 Gene
(In Gunn Rats)**

Vehicle	Dosage	G Clones/Total Clones	Frequency (%)
Gc-NLV	1 mg	112/815	15.4
		208/761	27.3
		185/974	18.9
		39/273	14.6 ¹
		78/403	19.3 ²
25 kDa PEI (Lactosylated)	1 mg	188/838	22.4
		254/1150	22.1
		245/997	24.6

¹Initial conversion frequency determined.

²Conversion frequency determined 7 days after 70% partial hepatectomy.

A Gunn rat was injected on five successive days with 1mg/Kg of CN3 complexed with 25 kDa Lac-PEI as above. Twenty five days after the final injection the serum bilirubin had declined from 6.2 mg/dl to 3.5 mg/dl and remained at that level for a further 25 days.

7.6 CORRECTION OF A FACTOR IX MUTATION IN DOG

The Chapel Hill strain of dogs, which has a (G→A)¹⁴⁷⁷ mutation that results in hemophilia in the animals, was used to obtain primary cultured hepatocytes. Four CMV to correct this mutation have been synthesized.

DIX1 (mut→WT) (SEQ ID No. 50)

```

T gcgcg auu caa aga aTT GAC CCT AAT AAT cga ccc cT
T                                     T
T                                     T
T CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG GT
      3' 5'

```

DIX2 (mut→WT) (SEQ ID No. 51)

```

T gcgcg caa aga auu gAC CCT AAT aau cga cT
T                                     T
T                                     T
T CGCGC GTT TCT TAA CTG GGA TTA TTA GCT GT
      3' 5'

```

DIX3 (mut→WT) (SEQ ID No. 52)

```

u gcgcg auu caa aga auu gac gcu aau aau cga ccc cu
u
u
u CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG Gu
      3' 5'

```

DIX4 (mut→WT) (SEQ ID No. 53)

```

u gcgcg auu caa aga auu gac ucu aau aau cga ccc cu
u
u
u CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG Gu
      3' 5'

```

DIX1 differs from DIX3 by the replacement of the intervening DNA segment with 2'-O-methyl RNA and replacement of the tetrathymidine linkers with tetrauracil. DIX 4 differs from DIX3 in that the mutational vector contains a mismatch in the mutator region. In DIX4 the 5' (lower) strand encodes the desired (wild-type) sequence while the 3' (upper) strand has the sequence of the target, i.e., the mutant sequence.

The hepatocytes were treated with 360 nM DIX1 complexed in either 25 kDa Lac-PEI or galactocerebroside-containing aqueous-cored, negatively charged lipid vesicles (Gc-NLV). The results are given in the table below.

Frequency of conversion of A to G at nucleotide 1477 of the Factor IX Gene

(Primary Hepatocytes from the Chapel Hill Strain of Hemophilia B Dogs)

Vehicle	Number of Times Transfected	Concentration	G Clones/Total Clones	Frequency (%)
Gc-NLV	Once	360 nM	30/195	15.44
			30/218	13.76
	Twice		30/118	25.4
Lac-PEI 25 kDa	Once*	360 nM	20/141	14.2
			48/348	13.3
	Twice		21/107	19.6

*RT-PCR on parallel transfected cultures gave an A to G conversion frequency of 11.1%

Each of the DIX2-DIX4 were also tested on primary cultured dog hepatocytes as above. The results showed that DIX2 worked poorly, possibly due to the low (25%) GC percentage. The subsequent experiments the results of DIX3 were about 16% conversion, while a parallel experiments DIX1 gave 10% conversion and the results of DIX4 were about as good as DIX1.

GenBank Sequence References for the Exons of the Human Apolipoprotein B-100 Gene

TABLE II

Exon No.	cDNA Boundary	GenBank Accession No. Sequence Reference
1	126 to 207	M19808
2	208 to 246	M19808
3	247 to 362	M19809
4	363 to 508	M19810
5	509 to 662	M19811
6	663 to 818	M19812
7	819 to 943	M19813
8	944 to 1029	M19813
9	1030 to 1249	M19815
10	1250 to 1477	M19816
11	1478 to 1595	M19818
12	1596 to 1742	M19818
13	1743 to 1954	M19820
14	1955 to 2192	M19820
15	2193 to 2359	M19821
16	2360 to 2561	M19823
17	2562 to 2729	M19824
18	2730 to 2941	M19824
19	2942 to 3124	M19825
20	3125 to 3246	M19825
21	3247 to 3457	M19827
22	3458 to 3633	M19828
23	3634 to 3821	M19828
24	3822 to 3967	M19828
25	3968 to 4341	M19828
26	4342 to 11913	M19828
27	11914 to 12028	M19828
28	12029 to 12212	M19828
29	12213 to 13816	M19828

SEQ ID	TABLE I			G/C#	NA Change	AA Change	AA	%APOB100	Restriction Site
No.	Sequence (5'→3')		Change						
4	AGTCTGGATGGG	I A A G C C C C C C T C A	15	A→T	K→Stop	1701	36.9	None	
5	CTGGGCTGGCTT	A A G C C A T T G A C A T	13	C→A	S→TAA	1876	40.8	+CTTAAG	
6	GCTCTCTGGGG	A A C A T A C T G G G C	14	G→T	E→Stop	1921	41.8	None	
7	GATGCCGTGAG	I A G C C C C C A A G A A T	13	A→T	K→Stop	2047	44.5	None	
8	GAGAGGAATCGA	I A A C C A T T A T A G	10	C→T	Q→Stop	2085	45.4	+ATCGAT	
9	TGTAAGAAAAA	T A A A G A G A G C G C C T G	10	C→A	Y→Stop	2110	45.9	None	
10	GCAGCCCTGGG	A I A A C T C C C A C A G C	16	A→T	K→Stop	2116	46.0	None	
11	GCAAGCTAATGATTAG	C T G A A T T C A A T T C A A T	8	T→G	Y→Stop	2124	46.2	+AGCT	
12	CAAGITCACATGCC	I A G G A G A A A C T G A C T G	11	A→T	K→Stop	2138	46.5	+CCTAGG	
13	ATATACAAAITGCA	I G A G A T G A T G C C A A A A T	9	T→G	L→Stop	2159	47.0	+CATG	
14	AAACTATCTCAACTG	I A G A C A T A T A T G A T A C	8	C→T	Q→Stop	2174	47.3	-CTGCCAG	
15	GCTAATATTATTCAT	I A A A T C A A T T G A A A A T T A	3	G→T	E→Stop	2204	48.0	+TTAA	
16	TGATGAGCACTAG	C A T A T C C G T G T A	11	T→G	Y→Stop	2216	48.3	+CTAG	
17	CTGCAGCAGCTT	I A G A G A C A C A T A C	12	A→T	K→Stop	2270	49.4	-CTTAAG	
18	AACAGTGAGCTG	I A G T G G C C C G I T C	14	C→T	Q→Stop	2684	58.6	None	
19	CAGACTCCGTT	A A C C A G A A A T C G C	12	T→A	L→Stop	2712	59.2	+GTTAAC	
20	AAAGGTCATGG	I A A T G G G C C T G C C	14	A→T	K→Stop	2930	64.0	None	
21	ACAATATGATA	I A A T T T G A T C A G T	5	C→T	Q→Stop	2180	47.5	Physiologic	

Table III

Seq ID No.	Sequence (5'→3')	G/C#	AA Change	NA Change	AA	Gene
22	ATGGAGGACGTGIGCGCGCCGCTGG	18	R→C	C→T	112	Apo E
23	GACCTGCAGAAAGIGCCCTGGCAGTGT	15	R→C	C→T	158	Apo E
24	GACCTGCAGAAAGCGCCCTGGCAGTGT	16	C→R	T→C	158	Apo E
25	TAAGGTCAGGAGITTGAGACCCAGCC	13	N/A	A→T	-491	Apo E
54	GCCGAGGACATGIGCGACCCGGCGGC	19	R→C	C→T	149	Apo AI
55	GAGATCGCGGACIGCGCGCGCGCGCC	20	R→C	C→T	151	Apo AI
56	CGCGACCGCGCGIGCGCGCATGTGG	20	R→C	C→T	153	Apo AI
57	AGCGACCAGCTGIGCCAGCGCCTGG	17	R→C	C→T	171	Apo AI
58	GAGCTGCGCCAGIGCTTGGCCCGCGC	19	R→C	C→T	173	Apo AI

WE CLAIM

1. A method of reducing LDL in the blood of a subject comprising altering an Apo B gene of a hepatocyte of the subject such that the transcript of the altered Apo B gene contains an in-frame stop codon whereby the altered gene encodes a protein having at least 1433 amino acids and not more than 3974 amino acids.
2. The method of claim 1, which further comprises the steps of determining the effect on the level of LDL of the alteration of the Apo B genes in the subject and subsequently adjusting the number of altered Apo B genes in the subject.
3. The method of claim 1, wherein the altered gene encodes a protein having at least 1841 amino acids and not more than 2975 amino acids.
4. The method of any one of claims 1-3, wherein the altered gene encodes a protein having a sequence of a fragment of SEQ ID No. 1, which fragment is at least amino acids 1-1841 and not more than amino acids 1-2975.
5. The method of any one of claims 1-4 which comprises administering a recombinagenic oligonucleobase which comprises a first and a second homologous region each having a sequence of at least 10 nucleobases selected from nt 4342-11913 of SEQ ID No: 2, whereby the alteration of the Apo B gene is effected.
6. The method of any one of claims 1-5, wherein the subject's fasting LDL serum cholesterol is reduced to below 140 mg/dl.
7. A composition for the modification of a human Apo B gene comprising an oligonucleobase which oligonucleobase comprises:
 - a. a first and a second homologous region that are each at least 8 nucleobases in length and together at least 20 nucleobases in length, which homologous regions are each homologous with a fragment of the sequence of nt 5649-9051 of SEQ ID No. 2, and

- b. a heterologous region that is disposed between the first and second homologous region,
such that the introduction of the sequence of the heterologous region into the Apo B gene results in the truncation of the protein encoded thereby.
- 8. The composition of claim 7, in which the first and the second homologous regions each comprises at least 3 contiguous nucleobase-pairs of hybrid-duplex.
- 9. The composition of claim 7 or 8, in which the sum of the lengths of the first and second homologous regions is not more than 60 nucleobases in length.
- 10. The composition of any one of claims 7-9, in which the homologous regions together comprise between 9 and 25 nucleobase pairs of hybrid-duplex.
- 11. The composition of any one of claims 7-10, in which the GC fraction of each homologous region is at least 33%.
- 12. The composition of any one of claims 7-10, in which the GC fraction of a homologous region is at least 50%.
- 13. The composition of any one of claims 7-10, in which the sequence of the oligonucleobase comprises the sequence of at least a 21 nucleobase fragment of any one of SEQ ID No. 4-21 or the complement thereof.
- 14. The composition of any one of claims 7-10, in which the sequence of the oligonucleobase comprises the sequence of at least a 25 nucleobase fragment of any one of SEQ ID No. 4-21 or the complement thereof.
- 15. The composition of any one of claims 7-14 which further comprises:
 - a. an aqueous carrier; and
 - b. a macromolecular carrier selected from the group consisting of

- i. an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - ii. a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - iii. a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
16. The composition of claim 15, in which the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
17. The composition of claim 16, in which the clathrin-coated pit receptor is selected from the group consisting of the receptors for transferrin, nicotinic acid, carnitine, insulin and insulin like growth factor-1.
18. The composition of claim 16, in which the clathrin-coated pit receptor is an asialoglycoprotein receptor.
19. The composition of any one of claims 15-18, in which the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
20. The composition of any one of claims 16-18, in which the aqueous-cored lipid vesicle further comprises a cerebroside.
21. The composition of any one of claims 15-20, in which the aqueous-cored lipid vesicle comprises dioleoylphosphatidylcholine and dioleoylphosphatidylserine.
22. The composition of any one of claims 16, 18, 19 and 21 wherein the macromolecular carrier is an aqueous-cored lipid vesicle that comprises a fusogenic F-protein.

23. The composition of any one of claims 15-22 , in which the oligonucleobase comprises:
- a. a first and a second homologous region that are together at least 16 and not more than 60 nucleobases in length, which regions are homologous with a target gene of a mammal; and
 - b. a heterologous region that is disposed between the first and second homologous region and is at least 1 and not more than 20 nucleobases in length, which is heterologous with the target gene and which contains the alteration.
24. A method of treatment and/or prophylaxis in a subject comprising altering an Apo E gene of a hepatocyte of the subject by introducing a substitution selected from the group (Arg→Cys)¹¹², (Arg→Cys)¹⁵⁸ and (Cys→Arg)¹⁵⁸.
25. The method of claim 24, wherein the subject is homozygous for Apo E4 and the alteration comprises the substitution (Arg→Cys)¹¹².
26. The method of claim 24 or 25, which comprises administering a chimeric mutational vector having a sequence which comprises SEQ ID No: 22.
27. The method of claim 24, wherein the treatment or prophylaxis comprises reducing the subject's fasting serum LDL cholesterol level and the alteration comprises the substitution (Arg→Cys)¹⁵⁸.
28. The method of claim 24 or 27, which comprises administering a chimeric mutational vector having a sequence which comprises SEQ ID No: 23.
29. The method of claim 24, wherein the subject suffers from Type III hyperlipidemia and the alteration comprises the substitution (Cys→Arg)¹⁵⁸.

30. The method of claim 24 or 29, which comprises administering a recombinagenic oligonucleobase having a sequence which comprises SEQ ID No: 24.
31. A composition for the alteration of a human Apo E gene comprising a recombinagenic oligonucleobase having a sequence comprising the sequence of at least a 21 nucleobase fragment of any one of SEQ ID No. 22 - 25 or the complement thereof.
32. A method of ameliorating atherosclerosis in a subject comprising altering an Apo A1 gene of a hepatocyte of the subject such that the altered Apo A1 protein forms dimers.
33. The method of claim 32, which further comprises the steps of determining the effect on the level of HDL of the alteration of the Apo A1 genes in the subject and subsequently adjusting the number of altered Apo A1 genes in the subject.
34. The method of claim 32, wherein the altered gene encodes a protein having a cysteine for arginine substitution at a position selected from the group consisting of residue 149, 151, 153, 171 and 173.
35. The method of claim 34, wherein the method comprises the administration of a recombinagenic oligonucleobase having a sequence comprising the sequence of at least 20 nucleotides of SEQ ID No. 54, No. 55, No. 56, No. 57 and No. 58.
36. The method of claim 35, which comprises the step of administering to the subject a composition comprising:
 - a) the recombinagenic oligonucleobase;
 - b) an aqueous carrier; and
 - c) a macromolecular carrier selected from the group consisting of

- (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
37. The method of claim 36, wherein the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
38. The method of claim 36, wherein the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
39. The method of claim 38, wherein the clathrin-coated pit receptor is an asialoglycoprotein receptor.
40. The method of claim 32, which comprises administering to the subject a composition comprising:
- a) a recombinagenic oligonucleobase;
 - b) an aqueous carrier; and
 - c) a macromolecular carrier selected from the group consisting of
 - (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
41. The method of claim 40, wherein the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.

42. The method of claim 40, wherein the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
43. The method of claim 42, wherein the clathrin-coated pit receptor is an asialoglycoprotein receptor.
44. A composition for the modification of a human Apo A1 gene comprising an oligonucleobase which oligonucleobase contains a sequence which comprises a fragment having the sequence selected from the group consisting of SEQ ID No. 54, No. 55, No. 56, No.57 and No. 58.
45. The composition of claim 44, which further comprises:
- a) an aqueous carrier; and
 - b) a macromolecular carrier selected from the group consisting of
 - (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
46. The composition of claim 45, in which the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
47. The composition of claim 46, in which the clathrin-coated pit receptor is selected from the group consisting of the receptors for transferrin, nicotinic acid, carnitine, insulin and insulin like growth factor-1.
48. The composition of claim 46, in which the clathrin-coated pit receptor is an asialoglycoprotein receptor.

49. The composition of any one of claims 45-48, in which the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
50. The composition of any one of claims 46-48, in which the aqueous-cored lipid vesicle further comprises a cerebroside.
51. The composition of any one of claims 45-50, in which the aqueous-cored lipid vesicle comprises dioleoylphosphatidylcholine and dioleoylphosphatidylserine.
52. The composition of any one of claims 46, 48, 49 and 51 wherein the macromolecular carrier is an aqueous-cored lipid vesicle that comprises a fusigenic F-protein.





FIG. 1

2/11

FIG. 2A

1 cctatccctgggggagggggcgggacaggggagccctataattggacaagtctgggatccttgagtcctACTCAGCCCCAG
83 CGGAGGTGAAGGACCTCTCCCCAGGAGCCGgtgagaagcgagtcgggggacggggatgagctcaggggcctctagaaa
165 gagctgggaccctgggaagccctggcctccaggtagctctcaggagagctactcggggtcgggcttggggagaggaggagcgg
247 ggggtgaggcaagcagcaggggactggacctgggaagggtcgggcagcagagacgacccgacccgctagaaggtgggggtgggg
329 agagcagctggactgggatgtaagccatagcaggactccacgagttgtcactatcattatcgagcacctactgggtgtcccc
411 agtgtcctcagatctccataactggggagccaggggcagcgacacggtagctagccgtcgattggagaactttaaaatgagg
493 actgaattagctcataaatggaacacggcgcttaactgtgaggttgagcttagaatgtgaagggagaatgaggaatgcgag
575 actgggactgagatggaaccggcggtggggaggggggtgggggatggaatttgaaacccgggagaggaagatggaattttct
657 atggaggccgacctggggatggggagataagagaagaccaggagggaggttaaatagggaatgggttggggcggtctggtaa
739 atgtgctgggattaggctgttcagataatgcaacaaggcttggaaggctaacctggggtgaggccgggttggggcgctgg
821 ggggtgggaggagtcctcactggcggttgattgacagtttctccttccccagACTGGCCAATCACAGGCAGGAAGATGAAGGT
M K V
903 TCTGTGGGCTGCGTTGCTGGTCACATTCCTGGCAGgtatggggcggggcttgctcgggttcccccgctccctccccctctca
L W A A L L V T F L A
985 tcctcacctcaacctcctggccccattcagacagacctgggccccctcttctgaggcttctgtgctgcttcctggctctga
1067 acagcgatttgacgctctctgggcctcggtttcccccatccttgagataggagtagaagttgtttgtgtgtgtgtgtgt
1149 tgttgtgtgtttgttttttgagatgaagtctcgctctgtcgccaggtggagtgagtgaggcgggatctcggtcactgca
1231 agctcgcctcccagggtccacgccattctcctgcctcagcctcccaagtagctgggactacaggcacatgccaccacacccg
1313 actaactttttgtattttcagtagagacgggggttcaccatgttggccaggctgggtctggaactcctgacctcaggtgatc

3/11

FIG. 2B

1395 tgcctcggttcgatctctccaaagtgtgtgggattacaggcgtgagccaccgcacctggctgggaggttagaggtttctaataatgcat
1477 tgcaggcagatagtgaaataccagacacggggcagctgtgatctttattctccatcacccccacacagccctgcctggggcac
1559 acaaggacactcaatacatgcttttccgctgggcccgtgtgtcaccctgtaatcccagcactttgggaggccaaggtggga
1641 ggatcacttgagcccaggagttcaacaccagcctgggcaacatagttagaccctgtctctactaaaaatacaaaaattagcc
1723 aggcattggtgccacacacctgtgtctctcagctactcaggaggctgaggcaggaggatcgcttgagcccagaaggtcaaggtt
1805 gcagtgaaccatgttcaggccgctgcactccagcctgggtgacagagcaagaccctgtttataatacataatgctttccaa
1887 gtgattaaaccgactccccctcacctgcccaccatgggtccaaagaagcattttgtggagcaccttctgtgtgccctagg
1969 tagctagatgcctggacggggtcagaaggaccctgaccgcacctgaacttgttccacacagGATGCCAGGCCAAGGTGGAG
G C Q A K V E
2051 CAAGCGGTGGAGACAGAGCCCGAGCCCGAGCTGCGCCAGCAGACCGAGTGGCAGAGCGGCCAGCGCTGGGAAGTGGCACTGG
Q A V E T E P E P E L R Q Q T E W Q S G Q R W E L A L
2133 GTCGCTTTTGGGATTACCTGCGCTGGGTGCAGACACTGTCTGAGCAGGTGCAGGAGGAGCTGCTCAGCTCCAGGTACCCCA
G R F W D Y L R W V Q T L S E Q V Q E E L L S S Q V T Q
2215 GGAAGTGAAGgtgagtgtccccatcctggcccttgaccctcctgggtgggcccgtatacctccccaggtccaggtttcattctg
E L R
2297 cccctgtcgctaagtcttggggggcctgggtctctgtgtgttctagcttccctcttccatttctgactcctggcttagctc
2379 tctggaattctctctctcagctttgtctctctctcttcccttctgactcagctctctcacactcgtcctggtctgtctctgt
2461 ccttccctagctcttttatatagagacagagagatgggggtctcactgtgttgcccaggctggtcttgaactctgggctcaa
2543 gcgatacctccgcctcgccctccaaagtgtgtgggattagaggcatgagcaccttgcccggcctcctagctccttctctgctc
2625 tctgctctgcccctctgcatctgtctctgcatctgtctctgtctccttctcteggcctctgcccgttccctctctcctc
2707 ttgggtctctctgtgtcatccccatctcggccgccccatcccagccttctccccgcctccccactgtgagacacccctccc

4/11

FIG. 2C

2789 gccctctcgggcgagggcgctgATGGACGAGACCATGAAGGAGTTGAAGGCCTACAAATCGGAACTGGAGGAACAACTGAC
A L M D E T M K E L K A Y K S E L E E Q L T

2871 CCGGTTGGCGGAGGAGACGCGGCGACGGCTGTCCAAGGAGCTGCAGGCGGCGCAGGCCCGGCTGGGCGCGGACATGGAGGAC
P V A E E T R A R L S K E L Q A A Q A R L G A D M E D

2953 GTGTGCGGCGCGCTGGTGCACTACCGCGGCGAGGTGCAGGCCATGCTCGGCCAGAGCACCGAGGAGCTGCGGGTGCGCCTCG
V C G R L V Q Y R G E V Q A M L G Q S T E E L R V R L

3035 CCTCCACCTGCGCAAGCTGCGTAAGCGGCTCCTCCGCGATGCGGATGACCTGCAGAAGCGCCTGGCAGTGTACCAGGCCGG
A S H L R K L R K R L L R D A D D L Q K R L A V Y Q A G

3117 GCGCGCGGAGGGGCGCGAGCGCGGCTCAGCGCCATCCGCGAGCGCCTCGGGCCCCCTGGTGGAACAGGGCGCGGTGGCGGGC
A R E G A E R G L S A I R E R L G P L V E Q G R V R A

3199 GCCACTGTGGGCTCCCTGGCGCGCCAGCGCTACAGGAGCGGGCCAGGCCCTGGGGCGAGCGGCTGCGCGCGGATGGAGG
A T V G S L A G Q P L Q E R A Q A W G E R L R A R M E

3281 AGATGGGCAGCCGACCGCGACCGCCTGGACGAGGTGAAGGAGCAGGTGGCGGAGGTGCGCGCAAGCTGGAGGAGCAGGC
E M G S R T R D R L D E V K E Q V A E V R A K L E E Q A

3363 CCAGCAGATACGCTGCAGGCGCGAGGCTTCCAGGCCCGCCTCAAGAGCTGGTTCGAGCCCTGGTGGAGACATGCAGCGC
Q Q I R L Q A E A F Q A R L K S W F E P L V E D M Q R

3445 CAGTGGGCGGGCTGGTGAGAAGGTGCAGGCTGCGGTGGGCACCGAGCGCGCCCTGTGCCAGCGACAATCACTGAACGC
Q W A G L V E K V Q A A V G 6 S A A P V P S D N H

3527 CGAAGCCTGCAGCCATGCGACCCACGCCACCCCGTGCCTCCTGCTCCGCGCAGCCTGCAGCGGAGACCTGTCCCGGCC

3609 CCAGCCGTCCTCCTGGGGTGGACCCCTAGTTTAATAAAGATTACCAAGTTTCAAGCAtctgtgtggcctccccctgtgatttc

3691 ctctaagccccagcctcagttttctctttctgtccacatactgcccacacaattctcagccccctctctccatctgtgtctgt

3373 gtgtatctttctctctgtgccccttttttttttttt

5/11

FIG. 3A

1 ctaaagaaga gcactggttg gaggacaggg cgggggaagg gggaggggag tgaagtagtc
 61 tccctggaat gctggtggtg ggggaggcag tctccttggg ggaggagtcc cagcgtccct
 121 cccctccct cctctgcaa cacaatggac ccaccccgag aatggcaact gccacacac tcccatggag
 181 gggaaagggg tgagtgcagg gaaccccgac cccacccggg agacctgcaa gcctgcagca
 241 cccccctcc cccccactg aaccttgac cctgcccctg cacgccccgc agcttgctgt
 301 ttgcccactc ctatttgccc agtcccagg acagagctga tccttgaact cttaaagtcc
 361 acattgccag gaccagtga gaccaacagg gccagggctg ggcttatcag cctccagcc
 421 cagaccctgg ctgcagacat aaataggccc tgcaagagct ggctgcttag agactgcgag
 481 aaggagggtg gtccctgctg ctgccccggc actctggctc ccagctcaa ggttcaggcc
 541 ttgccccagg ccgggcctct gggtaacctg ggtcttctcc cgctctgtgc ccttctctc
 601 acctggctgc aatgagtggg ggagcacggg gcttctgcat gctgaaggca cccactcag
 661 ccaggccctt cttctcctcc aggtccccca cggcccttca ggatgaaagc tgcggtgctg
 721 accttggccg tgctcttctt gacgggtagg tgtcccttaa cctaggagcc aaccatcggg
 781 gggcttcttc ctaaatccc cgtggcccac cctcctgggc agaggcagca ggttctcac
 841 tggccccctc tccccacct ccaagcttgg ccttctggct cagatctcag ccacagctg
 901 gcctgatctg ggtctccct cccacctca gggagccagg ctcgccattt ctggcagcaa
 961 gatgaacccc ccagagccc ctgggatcga tgaaaggacc tggccactgt gtacgtggat
 1021 gtgctcaaag acagcggcag agactatgtg tcccagttg aaggctccgc cttgggaaaa
 1081 cagctaaagt aaggaccag cctggggttg agggcagggg cagggggcag aggcctgtgg
 1141 gatgatgttg aagccagact ggcgagtcc tcacctata tctgatgagc tgggccccac
 1201 agatggtctg gatggagaaa ccggaatgga tctccaggca ggtcacagc ccatgtcccc
 1261 tgcaaaaggac agaccaggc tgcccgatgc gtgatcacag agccacattg tgcctgcaag
 1321 ttagcaagc cccttctcc tcttaccac cctctctgct cctgccccag aagactgtgg
 1381 gctgtcttcg gagaggagaa tgcgctggag gcatagaagc gaggtccttc aaggggccac

6/11

FIG. 3B

1441 tttagagacc aacgtaactg ggcaccagtc ccagctctgt ctcttttta gctcctctct
 1501 gtgcctcggc ccagctgcac aacggggcat ggcctggcgg ggcagggggtg ttggttgaga
 1561 gtgtactgga atgctaggc cactgcacct ccgcgacag gtgtcaccca gggctcaccc
 1621 ctgataggct gggcgctgg gaggccagcc ctcaaccctt ctgtctcacc ctccagccta
 1681 aagctccttg acaactggga cagcgtgacc tccaccttca gcaagctgcg cgaacagctc
 1741 ggcctctgtg cccaggagtt ctgggataac ctggaaaaagg agacagaggg cctgaggcaa
 1801 gagatgagca aggatctgga ggaggtgaag gccaaaggctg agccctacct ggacgacttc
 1861 cagaagaagt ggcaggagga gatggagctc taccgccaga aggtggagcc gctgcgcgca
 1921 gagctccaag agggcgcgcg ccagaagctg cacgagctgc aagagaagct gagccactg
 1981 ggcgaggaga tgcgcgacgg cgcgcgcgcc catgtggacg cgctgcgcac gcattctggc
 2041 ccctacagcg acgagctgcg ccagcgcttg gccgcgcgcc ttgaggctct caaggagaac
 2101 ggcggcgcca gactggccga gtaccacgcc aaggccaccg agcatctgag cagctcagc
 2161 gagaaggcca agcccgcgct cgaggacctc cgccaaggcc tgctgcccgt gctggagagc
 2221 ttcaaggcca gcttccctgag cgctctcgag ggtacacta agaagctcaa caccagtgga
 2281 ggcgcccgcg gccgcccccc ttcccggctg tcagaataaa cgtttccaaa gtgggaagca
 2341 gcttcttctt ttggggagaa tagagggggg tgccgggggaca tccggggggag cccgggaggg
 2401 gcctttggcc ctggagcagg gacttcctgc cggtatctca caactccgtg ccagactgg
 2461 acgtcttagg gccaaagatcg acgttgaggg acctgctgga cgcntggctg cttacgagtg
 2521 agggagtaga gtctgcctta gcaaggctca agtagaaagg aagtcacagc ggacnaggca
 2581 aagcdacaga caatccaaagg ccagggtgcc tgaaaggggc tcaaacaaagg cctgcagcct
 2641 gtctgagggc ggcaggaaa cagcggttgc tttagctggg agcagtcggg tccccgtcc
 2701 ccagaggtgt gtccgtatag agccttctcc agcccagccg ctgtcagcgg gcgggacgga
 2761 gcggggcgcc tcaggagacc agccactggg attgggggtt ggtcccgggt gcaagtgaag
 2821 cgcttggagt ttgcgcctgt cctcctttac taattcaaaa acctctcaaa cagacacttc

7/11

FIG. 3C

2881 ccttttcttc tcacaaggcc agtatccccc tcccactact cccatccccg ccagaacacg
 2941 ccgcggtctc ctcaggcaca gcagtggagg ccagtcctcc accccctgcg gctccatgcc
 3001 atgccacccc ctctttctgc cagccctggc agaagctggc ctgagtaaga aaattcacca
 3061 ccacctcttg caggtacatt ttattttcca agatgctctc atatctgtgc tctcactgca
 3121 tcctcccttc ccacatcct ggctagattg ccatacagc cagagcatgg atgaggacac
 3181 tgaagcctgg acctgtgacg tcgcttgccc agtgaacagc aggatgggct aggcagcgct
 3241 ttttagacc tgaccccctg gccatccatg attattgaaa agagtgtgcg ggtcgggtgc
 3301 ggtggctcaa gcctgtaatc ccagcacttt gggaggctga ggtgggcgta tcaactcagg
 3361 gccaggagtt tgagaccagc ctggccaata tggtgaaacc ctgtctctac taaaaataca
 3421 aaaaaaaat cagctggcat ggtggcttgc accgtaatc ccagctacta ggaaggctga
 3481 ggcaggagaa tcgcttgaa ctagggagga gagcaagact ccagctaaaa aaaaaaaag
 3541 gcactccagc ctgggcgacg ctgggagga aggtgcacag gagccgaaa tcatgccact
 3601 agtgtgtggc ctggcactca agttcacatg ggtgtgcagg catgcctgtg tattctcaca
 3661 tgacctccct gctcacggtc cctccttgca ctcatgtctg aatgtccccg ctgcacgcac
 3721 atggcttcac agatctggca gtgccttccc taccctctct ctgcaggggc ttttgcccc
 3781 tcatgcaggc ccctggataa tcggccccc cccatgtcc ccatctccag tgtatcttag
 3841 ctacctagg taaaggagtg ggcttttttag ttccctaacct tccagagcta caacagcagt
 3901 catccagcca ggtctgggtg ggaacatttt ctagatacgg gtgctgagat ctctcagccc
 3961 agagagaagc cctgggggat tttcagagag aaagcagtcct ccagggtggg ctggatgtac
 4021 tgatgccact gagatctgta aaggagtccc taacacctga cataggagtg acaaaactgt
 4081 tttctgcacc aactgagcag aatacacgca gctgacctgg gctcaaggct tggccctgcc
 4141 acgtgctggc tctgtgatgc tggccaagtg ccttcgcctc tccggggccac agttttttga
 4201 tctgaagagt ggagccctac tcaagccatc tgcagctctc gggctctctg acctgacatc
 4261 tttcgggttg tggggacaca aaggaagcag cctctattgg gagacctgt gcttcttttt

8/11

FIG. 3D

4321 ggtccccagga cactgcccc caccactcca gtccggtccc aagggtccag tcagctcaac
 4381 tgtaatcatg acaacattga tcaagcatct ttacgtgcag gtgctgtgcc aaacggttcg
 4441 aacgctctct catttcaatc tcacggcaaa cctacgggtg aggggtacg gttgtatcca
 4501 ctttacatgt aagaaactga ggtgatatac agtggtgga gccaaagaata gtgcctcgtt
 4561 gcatcttact ccaacctcta gccatccgg cctcctccct tcacgtgcgc ctaagagggc
 4621 taggtggcct ggatagggga ggtcagctcc acagttttga gtaaacacac acagtctcaa
 4681 ctctgatgac acttaagtgc caggcatagt ggctggcatg gggcacacac tcaagtcatg
 4741 ttgtgcagca cctaacagtt tatcaaatga tcagcaaaact tattgtcctg tttagacctc
 4801 cgcacaaagc tgtcaaggaa ggcaggggtac ggaggggtgat tcctacctta gagatgaaga
 4861 aactgaggcc cagagactag ccagctacc agaaggtgga tagagcgctg gcctccatgc
 4921 ctgcctgacc tggagtctgt ccagtgccta ccacagaac agcctcggcc ctttcccatg
 4981 ccagacaca gatggcacac ttgcgacggc ccactcatag cagcttcttg tccagcttta
 5041 ttgggaggcc agcatgcctg gagggggggc aggcattgagg tggggtaggga gaggactgag
 5101 aatactgtcc cttttaagca acctacaggg gaagccctgg agattgcagg acccaaggag
 5161 ctggcaggat ggataggcag gtggacttgg ggtattgagg tctcaggcag ccacggctga
 5221 agttggcttg acctcagggt ccaatccca gaactcagag aacttgtcct taacgggtgct
 5281 ccagtagtct tcagggaac tgaagccatc ggtcacccag cccctaaatc agtcagggga
 5341 agcaacagag cagggcata gaactcctct gtaggcaacc atgggaccca caccatgtc
 5401 cccactggac gacaccagtc aggaccacac caccctctca acttcactgg acgacagccc
 5461 tgagacctca ggcaggaatc cccacagct gccaccctgg gagaagagat atccttgag
 5521 gaaccccgag acaagtcaaa cctgacctc tccaggctac ctacagatgtt tatgccccg
 5581 ggcctgaggc acaaagtgc aggtggggga gatgtgaaag gtcaaggctg tcattgtttt
 5641 ctgtgcaaac agcaccacct ggagttgcac aacctgggtg ctctgagcag ggtaggacag
 5701 agggaggcag cctctcattt ggaaagtcat tggaggattg attagtgtg tgatctgggg

9/11

FIG. 3E

5761 caggtcacct aatcgcctctg agcctcaatt ttctcatctg caaagtgaga aaataacacc
 5821 taccccaaag cgggttcttg ggactaagaa tgtttatgaa cacctctgct atgccagcta
 5881 atgccagtca gggtagggtg gagaagggag taggggagag gagagtactg atgggcaggg
 5941 gcaggatggg agaggaagga cacacttctg ctcacactgg cccaggtct tagaacaag
 6001 cagaagcact cacgggcttg aatgggtca ggtggggcct cccaaggcaa acccctagcc
 6061 caggggtccc aaccccccaag tcatagatag gtactggccc atggactgtt aggagccggg
 6121 ctgcacagca cgaagtgagt ggtgggtgag tgagcaatac gcctgagctc cgctcctgt
 6181 cagatcaagt ggccttacat tctcatagga gtgtgaacct tattgtgaac tgcacatgcc
 6241 agggatctag gtggagggt ccttatgaga atctaatac tgatgatctg aggtggaaca
 6301 gcttcacccc aaaaccatcc cgcctggccc atggaaaaat tgtccaccac aaaactggtc
 6361 cctggtgcca aaagggtga ggaccactgc tctagccaga cttcagaaa aggaataatgg
 6421 ggccaggcgc agtggctcgt gcctgtaac ccagcacttt gggaggccga ggcaggagga
 6481 tcccctgagg tcaggaaatc aagaccaacc tggccaacat ggtgaaaccc catctctgct
 6541 aaaaatacaa aaattagctg ggtgtggttg cgcgtgcctg taatcccagc tacttgggag
 6601 gctgaggcag gagaatgggt tgaacccggg agacggagggt tgcagtgagc cgagatggca
 6661 cactgcact ccagcctggg tgacagaggg agactccatt aaaaaaaa gaagaagaaa
 6721 gaaagagaga aaggaaagaga ggaagagaaa gaaaaaggaa agaaagaaaag
 6781 aaagaaaaaga aagaaagaaa ggaagagaaa aaaggaaaag agaaagaaaaga
 6841 aagaaagaaa gaaagaaaaga aagaaaaata gaaagggaaa ggaagggaaa
 6901 atggaatatga gggctaaac ggcgcggccc taggactgct cggggagaaa ggagggacct
 6961 gggagggaatg cagtctcaac tctgtcatct ctccgcagc agcctgacaa aggccctgtg
 7021 aggaaagagg aggtgaaga gcacaggccc aggggagggt ggagcacctc cattccattg
 7081 ttgggatctc accagggcag gggcgggttg gaatggaggc agctggcagg ggggatggg
 7141 agggaggcca gcgggtgtac ctggcctgct gggccacctg ggactcctgc acgctgctca

10/11

FIG. 3F

7201 gtgcacccctt ggcggtcttg gtggcgctgct tcatgtagcc ctgcatgaag ctgagaaggg
7261 aggcacccctc ggcctctgaa gctcctgagg aaagagcagg gctgagtggg gtggatcggc
7321 ctctggacga gccctggggc tcctgcttga ccaccattg gactgggac ccgaagtgc
7381 ctccaccctg cccccagccc agtcccacca agtgcttacg ggcagagggc aggagcgcca
7441 ggagggaac aacaaggagt acccggggct gcatggcacc tctgttcctg caaggaagtg
7501 tcctgtgagg ggcacccccc gtcaccatgc ctctggaccc ctccctgggg aggtggcgtg
7561 gcccttaagg tagaacctta gctgggtctg ccagaaggag taggggcccg ctctgctca
7621 tacgggctct cagaaggggg actggtgagc ggcgagggat cgaggcccaa aggaggtggg
7681 tgggatggag cagaaaacc accagactga acatcaaggc actgcggtct ggactgatct
7741 ccgtccagtc cagccaacat gctgtgtgtc ttgggtgat ttctggccct ctccaggcc
7801 tcagtttccc tgtctgggt aggaactggc tgtctaaggc ttctccagc cctaagcctg
7861 aagaatgagg ggggaacctg cacttgagc cacttccagc cccacccct atgtagcttt
7921 gggcaagtga caccctcccg gcctccatgt tcttcagggt atgatgagg gtggggggca
7981 cccgtccagc tccgaggctt ccttagctct agcaagtgtc ttctccaggc ttctggctgg
8041 gctgggcagg gagctcctct tgccctctct catcctcctc cctcctctt tccctcccc
8101 agagggcatt acctggagca gctgcctcta gggatgaact ggcagagacg gcaggaggg
8161 tctgacctgt ttatatcat ctccaggga caggcactg aggacccagg gcgctgggca
8221 aaggcacct gctgaccagt ggagatgagg gcctgaggca ggggtgtccag atgcagcaag
8281 cgggcgggag agttgggaaa tccctaggag actgagtcca cgctgctgtc ccgccagccc
8341 tgacgcccag atgagctcag gaactggggg tgggcctggg gacctcatt ccttccctagc
8401 tgactggctc ccagggaga ggctgggtgag aggggaaatg gcttgacat agggcagcgc
8461 tcgggcccac ctacgcttt cactctggaa ttctaggccc ctccctccac cagccccaa
8521 ccctgaacac agcctggagt agagggtga ggggcttctt cagacttgag acaagtggg
8581 tggcttgggc tggggggtgt ttggagtaaa ggcacagaag accgggcac agtggctcgc

11/11

FIG. 3G

8641 aaggccctggt tacagggctg agtcacagc ccctcccagc acctccatct ctgggtttca
8701 atccaggcag gcgagtgtg ctcacccagc agccccccac ccgccccac cctgtgtgcc
8761 ccaccgccgc ctccccctga gtgtaggcca ggggttggtg gagaagcgca aggccgctca
8821 gagccccgagg cctttgcccc tccctccacc aggtcccta tttgcccctc tggaccact
8881 gataaacatcc ctgggggagg agccccacc aggaactgatt ctctcgttca ccttcctggg
8941 ttcctcagtg gggctggggg aagagc

SEQUENCE LISTING

<110> 1. Steer, Clifford J.
 2. Kren, Betsy T.
 3. Bandyopadhyay, Paramita
 4. Roy-Chowdhury, Jayanta

<120> Methods and Compounds for the Genetic
 Treatment of Hyperlipidemia

<130> 7991-033-228

<150> 60/074,497

<151> 1998-02-12

<150> 09/108,006

<151> 1998-06-30

<160> 59

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4563

<212> PRT

<213> Homo Sapiens

<400> 1

```

Met Asp Pro Pro Arg Pro Ala Leu Leu Ala Leu Leu Ala Leu Pro Ala
 1              5              10              15
Leu Leu Leu Leu Leu Leu Ala Gly Ala Arg Ala Glu Glu Glu Met Leu
      20              25              30
Glu Asn Val Ser Leu Val Cys Pro Lys Asp Ala Thr Arg Phe Lys His
      35              40              45
Leu Arg Lys Tyr Thr Tyr Asn Tyr Glu Ala Glu Ser Ser Ser Gly Val
      50              55              60
Pro Gly Thr Ala Asp Ser Arg Ser Ala Thr Arg Ile Asn Cys Lys Val
      65              70              75              80
Glu Leu Glu Val Pro Gln Leu Cys Ser Phe Ile Leu Lys Thr Ser Gln
      85              90              95
Cys Ile Leu Lys Glu Val Tyr Gly Phe Asn Pro Glu Gly Lys Ala Leu
      100             105             110
Leu Lys Lys Thr Lys Asn Ser Glu Glu Phe Ala Ala Ala Met Ser Arg
      115             120             125
Tyr Glu Leu Lys Leu Ala Ile Pro Glu Gly Lys Gln Val Phe Leu Tyr
      130             135             140
Pro Glu Lys Asp Glu Pro Thr Tyr Ile Leu Asn Ile Lys Arg Gly Ile
      145             150             155             160
Ile Ser Ala Leu Leu Val Pro Pro Glu Thr Glu Glu Ala Lys Gln Val
      165             170             175
Leu Phe Leu Asp Thr Val Tyr Gly Asn Cys Ser Thr His Phe Thr Val
      180             185             190
Lys Thr Arg Lys Gly Asn Val Ala Thr Glu Ile Ser Thr Glu Arg Asp

```

Lys Ser Val Ser Ile Pro Ser Leu Asp Pro Ala Ser Ala Lys Ile Glu
 645 650 655
 Gly Asn Leu Ile Phe Asp Pro Asn Asn Tyr Leu Pro Lys Glu Ser Met
 660 665 670
 Leu Lys Thr Thr Leu Thr Ala Phe Gly Phe Ala Ser Ala Asp Leu Ile
 675 680 685
 Glu Ile Gly Leu Glu Gly Lys Gly Phe Glu Pro Thr Leu Glu Ala Leu
 690 695 700
 Phe Gly Lys Gln Gly Phe Phe Pro Asp Ser Val Asn Lys Ala Leu Tyr
 705 710 715 720
 Trp Val Asn Gly Gln Val Pro Asp Gly Val Ser Lys Val Leu Val Asp
 725 730 735
 His Phe Gly Tyr Thr Lys Asp Asp Lys His Glu Gln Asp Met Val Asn
 740 745 750
 Gly Ile Met Leu Ser Val Glu Lys Leu Ile Lys Asp Leu Lys Ser Lys
 755 760 765
 Glu Val Pro Glu Ala Arg Ala Tyr Leu Arg Ile Leu Gly Glu Glu Leu
 770 775 780
 Gly Phe Ala Ser Leu His Asp Leu Gln Leu Leu Gly Lys Leu Leu Leu
 785 790 795 800
 Met Gly Ala Arg Thr Leu Gln Gly Ile Pro Gln Met Ile Gly Glu Val
 805 810 815
 Ile Arg Lys Gly Ser Lys Asn Asp Phe Phe Leu His Tyr Ile Phe Met
 820 825 830
 Glu Asn Ala Phe Glu Leu Pro Thr Gly Ala Gly Leu Gln Leu Gln Ile
 835 840 845
 Ser Ser Ser Gly Val Ile Ala Pro Gly Ala Lys Ala Gly Val Lys Leu
 850 855 860
 Glu Val Ala Asn Met Gln Ala Glu Leu Val Ala Lys Pro Ser Val Ser
 865 870 875 880
 Val Glu Phe Val Thr Asn Met Gly Ile Ile Ile Pro Asp Phe Ala Arg
 885 890 895
 Ser Gly Val Gln Met Asn Thr Asn Phe Phe His Glu Ser Gly Leu Glu
 900 905 910
 Ala His Val Ala Leu Lys Pro Gly Lys Leu Lys Phe Ile Ile Pro Ser
 915 920 925
 Pro Lys Arg Pro Val Lys Leu Leu Ser Gly Gly Asn Thr Leu His Leu
 930 935 940
 Val Ser Thr Thr Lys Thr Glu Val Ile Pro Pro Leu Ile Glu Asn Arg
 945 950 955 960
 Gln Ser Trp Ser Val Cys Lys Gln Val Phe Pro Gly Leu Asn Tyr Cys
 965 970 975
 Thr Ser Gly Ala Tyr Ser Asn Ala Ser Ser Thr Asp Ser Ala Ser Tyr
 980 985 990
 Tyr Pro Leu Thr Gly Asp Thr Arg Leu Glu Leu Glu Leu Arg Pro Thr
 995 1000 1005
 Gly Glu Ile Glu Gln Tyr Ser Val Ser Ala Thr Tyr Glu Leu Gln Arg
 1010 1015 1020
 Glu Asp Arg Ala Leu Val Asp Thr Leu Lys Phe Val Thr Gln Ala Glu
 1025 1030 1035 1040
 Gly Ala Lys Gln Thr Glu Ala Thr Met Thr Phe Lys Tyr Asn Arg Gln
 1045 1050 1055
 Ser Met Thr Leu Ser Ser Glu Val Gln Ile Pro Asp Phe Asp Val Asp
 1060 1065 1070
 Leu Gly Thr Ile Leu Arg Val Asn Asp Glu Ser Thr Glu Gly Lys Thr

1075	1080	1085
Ser Tyr Arg Leu Thr Leu Asp Ile Gln Asn Lys Lys Ile Thr Glu Val		
1090	1095	1100
Ala Leu Met Gly His Leu Ser Cys Asp Thr Lys Glu Glu Arg Lys Ile		
1105	1110	1115
Lys Gly Val Ile Ser Ile Pro Arg Leu Gln Ala Glu Ala Arg Ser Glu		
1125	1130	1135
Ile Leu Ala His Trp Ser Pro Ala Lys Leu Leu Leu Gln Met Asp Ser		
1140	1145	1150
Ser Ala Thr Ala Tyr Gly Ser Thr Val Ser Lys Arg Val Ala Trp His		
1155	1160	1165
Tyr Asp Glu Glu Lys Ile Glu Phe Glu Trp Asn Thr Gly Thr Asn Val		
1170	1175	1180
Asp Thr Lys Lys Met Thr Ser Asn Phe Pro Val Asp Leu Ser Asp Tyr		
1185	1190	1195
Pro Lys Ser Leu His Met Tyr Ala Asn Arg Leu Leu Asp His Arg Val		
1205	1210	1215
Pro Gln Thr Asp Met Thr Phe Arg His Val Gly Ser Lys Leu Ile Val		
1220	1225	1230
Ala Met Ser Ser Trp Leu Gln Lys Ala Ser Gly Ser Leu Pro Tyr Thr		
1235	1240	1245
Gln Thr Leu Gln Asp His Leu Asn Ser Leu Lys Glu Phe Asn Leu Gln		
1250	1255	1260
Asn Met Gly Leu Pro Asp Phe His Ile Pro Glu Asn Leu Phe Leu Lys		
1265	1270	1275
Ser Asp Gly Arg Val Lys Tyr Thr Leu Asn Lys Asn Ser Leu Lys Ile		
1285	1290	1295
Glu Ile Pro Leu Pro Phe Gly Gly Lys Ser Ser Arg Asp Leu Lys Met		
1300	1305	1310
Leu Glu Thr Val Arg Thr Pro Ala Leu His Phe Lys Ser Val Gly Phe		
1315	1320	1325
His Leu Pro Ser Arg Glu Phe Gln Val Pro Thr Phe Thr Ile Pro Lys		
1330	1335	1340
Leu Tyr Gln Leu Gln Val Pro Leu Leu Gly Val Leu Asp Leu Ser Thr		
1345	1350	1355
Asn Val Tyr Ser Asn Leu Tyr Asn Trp Ser Ala Ser Tyr Ser Gly Gly		
1365	1370	1375
Asn Thr Ser Thr Asp His Phe Ser Leu Arg Ala Arg Tyr His Met Lys		
1380	1385	1390
Ala Asp Ser Val Val Asp Leu Leu Ser Tyr Asn Val Gln Gly Ser Gly		
1395	1400	1405
Glu Thr Thr Tyr Asp His Lys Asn Thr Phe Thr Leu Ser Cys Asp Gly		
1410	1415	1420
Ser Leu Arg His Lys Phe Leu Asp Ser Asn Ile Lys Phe Ser His Val		
1425	1430	1435
Glu Lys Leu Gly Asn Asn Pro Val Ser Lys Gly Leu Leu Ile Phe Asp		
1445	1450	1455
Ala Ser Ser Ser Trp Gly Pro Gln Met Ser Ala Ser Val His Leu Asp		
1460	1465	1470
Ser Lys Lys Lys Gln His Leu Phe Val Lys Glu Val Lys Ile Asp Gly		
1475	1480	1485
Gln Phe Arg Val Ser Ser Phe Tyr Ala Lys Gly Thr Tyr Gly Leu Ser		
1490	1495	1500
Cys Gln Arg Asp Pro Asn Thr Gly Arg Leu Asn Gly Glu Ser Asn Leu		
1505	1510	1515
		1520

Arg Phe Asn Ser Ser Tyr Leu Gln Gly Thr Asn Gln Ile Thr Gly Arg
 1525 1530 1535
 Tyr Glu Asp Gly Thr Leu Ser Leu Thr Ser Thr Ser Asp Leu Gln Ser
 1540 1545 1550
 Gly Ile Ile Lys Asn Thr Ala Ser Leu Lys Tyr Glu Asn Tyr Glu Leu
 1555 1560 1565
 Thr Leu Lys Ser Asp Thr Asn Gly Lys Tyr Lys Asn Phe Ala Thr Ser
 1570 1575 1580
 Asn Lys Met Asp Met Thr Phe Ser Lys Gln Asn Ala Leu Leu Arg Ser
 1585 1590 1595 1600
 Glu Tyr Gln Ala Asp Tyr Glu Ser Leu Arg Phe Phe Ser Leu Leu Ser
 1605 1610 1615
 Gly Ser Leu Asn Ser His Gly Leu Glu Leu Asn Ala Asp Ile Leu Gly
 1620 1625 1630
 Thr Asp Lys Ile Asn Ser Gly Ala His Lys Ala Thr Leu Arg Ile Gly
 1635 1640 1645
 Gln Asp Gly Ile Ser Thr Ser Ala Thr Thr Asn Leu Lys Cys Ser Leu
 1650 1655 1660
 Leu Val Leu Glu Asn Glu Leu Asn Ala Glu Leu Gly Leu Ser Gly Ala
 1665 1670 1675 1680
 Ser Met Lys Leu Thr Thr Asn Gly Arg Phe Arg Glu His Asn Ala Lys
 1685 1690 1695
 Phe Ser Leu Asp Gly Lys Ala Ala Leu Thr Glu Leu Ser Leu Gly Ser
 1700 1705 1710
 Ala Tyr Gln Ala Met Ile Leu Gly Val Asp Ser Lys Asn Ile Phe Asn
 1715 1720 1725
 Phe Lys Val Ser Gln Glu Gly Leu Lys Leu Ser Asn Asp Met Met Gly
 1730 1735 1740
 Ser Tyr Ala Glu Met Lys Phe Asp His Thr Asn Ser Leu Asn Ile Ala
 1745 1750 1755 1760
 Gly Leu Ser Leu Asp Phe Ser Ser Lys Leu Asp Asn Ile Tyr Ser Ser
 1765 1770 1775
 Asp Lys Phe Tyr Lys Gln Thr Val Asn Leu Gln Leu Gln Pro Tyr Ser
 1780 1785 1790
 Leu Val Thr Thr Leu Asn Ser Asp Leu Lys Tyr Asn Ala Leu Asp Leu
 1795 1800 1805
 Thr Asn Asn Gly Lys Leu Arg Leu Glu Pro Leu Lys Leu His Val Ala
 1810 1815 1820
 Gly Asn Leu Lys Gly Ala Tyr Gln Asn Asn Glu Ile Lys His Ile Tyr
 1825 1830 1835 1840
 Ala Ile Ser Ser Ala Ala Leu Ser Ala Ser Tyr Lys Ala Asp Thr Val
 1845 1850 1855
 Ala Lys Val Gln Gly Val Glu Phe Ser His Arg Leu Asn Thr Asp Ile
 1860 1865 1870
 Ala Gly Leu Ala Ser Ala Ile Asp Met Ser Thr Asn Tyr Asn Ser Asp
 1875 1880 1885
 Ser Leu His Phe Ser Asn Val Phe Arg Ser Val Met Ala Pro Phe Thr
 1890 1895 1900
 Met Thr Ile Asp Ala His Thr Asn Gly Asn Gly Lys Leu Ala Leu Trp
 1905 1910 1915 1920
 Gly Glu His Thr Gly Gln Leu Tyr Ser Lys Phe Leu Leu Lys Ala Glu
 1925 1930 1935
 Pro Leu Ala Phe Thr Phe Ser His Asp Tyr Lys Gly Ser Thr Ser His
 1940 1945 1950
 His Leu Val Ser Arg Lys Ser Ile Ser Ala Ala Leu Glu His Lys Val

1955	1960	1965
Ser Ala Leu Leu Thr Pro	Ala Glu Gln Thr Gly Thr Trp Lys Leu Lys	
1970	1975	1980
Thr Gln Phe Asn Asn Asn Glu Tyr Ser Gln Asp Leu Asp Ala Tyr Asn		
1985	1990	1995
Thr Lys Asp Lys Ile Gly Val Glu Leu Thr Gly Arg Thr Leu Ala Asp		2000
2005	2010	2015
Leu Thr Leu Leu Asp Ser Pro Ile Lys Val Pro Leu Leu Leu Ser Glu		
2020	2025	2030
Pro Ile Asn Ile Ile Asp Ala Leu Glu Met Arg Asp Ala Val Glu Lys		
2035	2040	2045
Pro Gln Glu Phe Thr Ile Val Ala Phe Val Lys Tyr Asp Lys Asn Gln		
2050	2055	2060
Asp Val His Ser Ile Asn Leu Pro Phe Phe Glu Thr Leu Gln Glu Tyr		
2065	2070	2075
Phe Glu Arg Asn Arg Gln Thr Ile Ile Val Val Leu Glu Asn Val Gln		2080
2085	2090	2095
Arg Asn Leu Lys His Ile Asn Ile Asp Gln Phe Val Arg Lys Tyr Arg		
2100	2105	2110
Ala Ala Leu Gly Lys Leu Pro Gln Gln Ala Asn Asp Tyr Leu Asn Ser		
2115	2120	2125
Phe Asn Trp Glu Arg Gln Val Ser His Ala Lys Glu Lys Leu Thr Ala		
2130	2135	2140
Leu Thr Lys Lys Tyr Arg Ile Thr Glu Asn Asp Ile Gln Ile Ala Leu		
2145	2150	2155
Asp Asp Ala Lys Ile Asn Phe Asn Glu Lys Leu Ser Gln Leu Gln Thr		
2165	2170	2175
Tyr Met Ile Gln Phe Asp Gln Tyr Ile Lys Asp Ser Tyr Asp Leu His		
2180	2185	2190
Asp Leu Lys Ile Ala Ile Ala Asn Ile Ile Asp Glu Ile Ile Glu Lys		
2195	2200	2205
Leu Lys Ser Leu Asp Glu His Tyr His Ile Arg Val Asn Leu Val Lys		
2210	2215	2220
Thr Ile His Asp Leu His Leu Phe Ile Glu Asn Ile Asp Phe Asn Lys		
2225	2230	2235
Ser Gly Ser Ser Thr Ala Ser Trp Ile Gln Asn Val Asp Thr Lys Tyr		2240
2245	2250	2255
Gln Ile Arg Ile Gln Ile Gln Glu Lys Leu Gln Gln Leu Lys Arg His		
2260	2265	2270
Ile Gln Asn Ile Asp Ile Gln His Leu Ala Gly Lys Leu Lys Gln His		
2275	2280	2285
Ile Glu Ala Ile Asp Val Arg Val Leu Leu Asp Gln Leu Gly Thr Thr		
2290	2295	2300
Ile Ser Phe Glu Arg Ile Asn Asp Val Leu Glu His Val Lys His Phe		
2305	2310	2315
Val Ile Asn Leu Ile Gly Asp Phe Glu Val Ala Glu Lys Ile Asn Ala		2320
2325	2330	2335
Phe Arg Ala Lys Val His Glu Leu Ile Glu Arg Tyr Glu Val Asp Gln		
2340	2345	2350
Gln Ile Gln Val Leu Met Asp Lys Leu Val Glu Leu Ala His Gln Tyr		
2355	2360	2365
Lys Leu Lys Glu Thr Ile Gln Lys Leu Ser Asn Val Leu Gln Gln Val		
2370	2375	2380
Lys Ile Lys Asp Tyr Phe Glu Lys Leu Val Gly Phe Ile Asp Asp Ala		
2385	2390	2395
		2400

Val Lys Lys Leu Asn Glu Leu Ser Phe Lys Thr Phe Ile Glu Asp Val
 2405 2410 2415
 Asn Lys Phe Leu Asp Met Leu Ile Lys Lys Leu Lys Ser Phe Asp Tyr
 2420 2425 2430
 His Gln Phe Val Asp Glu Thr Asn Asp Lys Ile Arg Glu Val Thr Gln
 2435 2440 2445
 Arg Leu Asn Gly Glu Ile Gln Ala Leu Glu Leu Pro Gln Lys Ala Glu
 2450 2455 2460
 Ala Leu Lys Leu Phe Leu Glu Glu Thr Lys Ala Thr Val Ala Val Tyr
 2465 2470 2475 2480
 Leu Glu Ser Leu Gln Asp Thr Lys Ile Thr Leu Ile Ile Asn Trp Leu
 2485 2490 2495
 Gln Glu Ala Leu Ser Ser Ala Ser Leu Ala His Met Lys Ala Lys Phe
 2500 2505 2510
 Arg Glu Thr Leu Glu Asp Thr Arg Asp Arg Met Tyr Gln Met Asp Ile
 2515 2520 2525
 Gln Gln Glu Leu Gln Arg Tyr Leu Ser Leu Val Gly Gln Val Tyr Ser
 2530 2535 2540
 Thr Leu Val Thr Tyr Ile Ser Asp Trp Trp Thr Leu Ala Ala Lys Asn
 2545 2550 2555 2560
 Leu Thr Asp Phe Ala Glu Gln Tyr Ser Ile Gln Asp Trp Ala Lys Arg
 2565 2570 2575
 Met Lys Ala Leu Val Glu Gln Gly Phe Thr Val Pro Glu Ile Lys Thr
 2580 2585 2590
 Ile Leu Gly Thr Met Pro Ala Phe Glu Val Ser Leu Gln Ala Leu Gln
 2595 2600 2605
 Lys Ala Thr Phe Gln Thr Pro Asp Phe Ile Val Pro Leu Thr Asp Leu
 2610 2615 2620
 Arg Ile Pro Ser Val Gln Ile Asn Phe Lys Asp Leu Lys Asn Ile Lys
 2625 2630 2635 2640
 Ile Pro Ser Arg Phe Ser Thr Pro Glu Phe Thr Ile Leu Asn Thr Phe
 2645 2650 2655
 His Ile Pro Ser Phe Thr Ile Asp Phe Val Glu Met Lys Val Lys Ile
 2660 2665 2670
 Ile Arg Thr Ile Asp Gln Met Leu Asn Ser Glu Leu Gln Trp Pro Val
 2675 2680 2685
 Pro Asp Ile Tyr Leu Arg Asp Leu Lys Val Glu Asp Ile Pro Leu Ala
 2690 2695 2700
 Arg Ile Thr Leu Pro Asp Phe Arg Leu Pro Glu Ile Ala Ile Pro Glu
 2705 2710 2715 2720
 Phe Ile Ile Pro Thr Leu Asn Leu Asn Asp Phe Gln Val Pro Asp Leu
 2725 2730 2735
 His Ile Pro Glu Phe Gln Leu Pro His Ile Ser His Thr Ile Glu Val
 2740 2745 2750
 Pro Thr Phe Gly Lys Leu Tyr Ser Ile Leu Lys Ile Gln Ser Pro Leu
 2755 2760 2765
 Phe Thr Leu Asp Ala Asn Ala Asp Ile Gly Asn Gly Thr Thr Ser Ala
 2770 2775 2780
 Asn Glu Ala Gly Ile Ala Ala Ser Ile Thr Ala Lys Gly Glu Ser Lys
 2785 2790 2795 2800
 Leu Glu Val Leu Asn Phe Asp Phe Gln Ala Asn Ala Gln Leu Ser Asn
 2805 2810 2815
 Pro Lys Ile Asn Pro Leu Ala Leu Lys Glu Ser Val Lys Phe Ser Ser
 2820 2825 2830
 Lys Tyr Leu Arg Thr Glu His Gly Ser Glu Met Leu Phe Phe Gly Asn

2835	2840	2845
Ala Ile Glu Gly Lys Ser Asn Thr Val	Ala Ser Leu His Thr Glu Lys	
2850	2855	2860
Asn Thr Leu Glu Leu Ser Asn Gly Val	Ile Val Lys Ile Asn Asn Gln	
2865	2870	2875
Leu Thr Leu Asp Ser Asn Thr Lys Tyr Phe	His Lys Leu Asn Ile Pro	2880
2885	2890	2895
Lys Leu Asp Phe Ser Ser Gln Ala Asp	Leu Arg Asn Glu Ile Lys Thr	
2900	2905	2910
Leu Leu Lys Ala Gly His Ile Ala Trp	Thr Ser Ser Gly Lys Gly Ser	
2915	2920	2925
Trp Lys Trp Ala Cys Pro Arg Phe Ser	Asp Glu Gly Thr His Glu Ser	
2930	2935	2940
Gln Ile Ser Phe Thr Ile Glu Gly Pro	Leu Thr Ser Phe Gly Leu Ser	
2945	2950	2955
Asn Lys Ile Asn Ser Lys His Leu Arg	Val Asn Gln Asn Leu Val Tyr	2960
2965	2970	2975
Glu Ser Gly Ser Leu Asn Phe Ser Lys	Leu Glu Ile Gln Ser Gln Val	
2980	2985	2990
Asp Ser Gln His Val Gly His Ser Val	Leu Thr Ala Lys Gly Met Ala	
2995	3000	3005
Leu Phe Gly Glu Gly Lys Ala Glu Phe	Thr Gly Arg His Asp Ala His	
3010	3015	3020
Leu Asn Gly Lys Val Ile Gly Thr Leu	Lys Asn Ser Leu Phe Phe Ser	
3025	3030	3035
Ala Gln Pro Phe Glu Ile Thr Ala Ser	Thr Asn Asn Glu Gly Asn Leu	3040
3045	3050	3055
Lys Val Arg Phe Pro Leu Arg Leu Thr	Gly Lys Ile Asp Phe Leu Asn	
3060	3065	3070
Asn Tyr Ala Leu Phe Leu Ser Pro Ser	Ala Gln Gln Ala Ser Trp Gln	
3075	3080	3085
Val Ser Ala Arg Phe Asn Gln Tyr Lys	Tyr Asn Gln Asn Phe Ser Ala	
3090	3095	3100
Gly Asn Asn Glu Asn Ile Met Glu Ala	His Val Gly Ile Asn Gly Glu	
3105	3110	3115
Ala Asn Leu Asp Phe Leu Asn Ile Pro	Leu Thr Ile Pro Glu Met Arg	
3125	3130	3135
Leu Pro Tyr Thr Ile Ile Thr Thr Pro	Pro Pro Leu Lys Asp Phe Ser Leu	
3140	3145	3150
Trp Glu Lys Thr Gly Leu Lys Glu Phe	Leu Lys Thr Thr Lys Gln Ser	
3155	3160	3165
Phe Asp Leu Ser Val Lys Ala Gln Tyr	Lys Lys Asn Lys His Arg His	
3170	3175	3180
Ser Ile Thr Asn Pro Leu Ala Val Leu	Cys Glu Phe Ile Ser Gln Ser	
3185	3190	3195
Ile Lys Ser Phe Asp Arg His Phe Glu	Lys Asn Arg Asn Asn Ala Leu	
3205	3210	3215
Asp Phe Val Thr Lys Ser Tyr Asn Glu	Thr Lys Ile Lys Phe Asp Lys	
3220	3225	3230
Tyr Lys Ala Glu Lys Ser His Asp Glu	Leu Pro Arg Thr Phe Gln Ile	
3235	3240	3245
Pro Gly Tyr Thr Val Pro Val Val Asn	Val Glu Val Ser Pro Phe Thr	
3250	3255	3260
Ile Glu Met Ser Ala Phe Gly Tyr Val	Phe Pro Lys Ala Val Ser Met	
3265	3270	3275
		3280

Pro Ser Phe Ser Ile Leu Gly Ser Asp Val Arg Val Pro Ser Tyr Thr
 3285 3290 3295
 Leu Ile Leu Pro Ser Leu Glu Leu Pro Val Leu His Val Pro Arg Asn
 3300 3305 3310
 Leu Lys Leu Ser Leu Pro Asp Phe Lys Glu Leu Cys Thr Ile Ser His
 3315 3320 3325
 Ile Phe Ile Pro Ala Met Gly Asn Ile Thr Tyr Asp Phe Ser Phe Lys
 3330 3335 3340
 Ser Ser Val Ile Thr Leu Asn Thr Asn Ala Glu Leu Phe Asn Gln Ser
 3345 3350 3355 3360
 Asp Ile Val Ala His Leu Leu Ser Ser Ser Ser Ser Val Ile Asp Ala
 3365 3370 3375
 Leu Gln Tyr Lys Leu Glu Gly Thr Thr Arg Leu Thr Arg Lys Arg Gly
 3380 3385 3390
 Leu Lys Leu Ala Thr Ala Leu Ser Leu Ser Asn Lys Phe Val Glu Gly
 3395 3400 3405
 Ser His Asn Ser Thr Val Ser Leu Thr Thr Lys Asn Met Glu Val Ser
 3410 3415 3420
 Val Ala Thr Thr Thr Lys Ala Gln Ile Pro Ile Leu Arg Met Asn Phe
 3425 3430 3435 3440
 Lys Gln Glu Leu Asn Gly Asn Thr Lys Ser Lys Pro Thr Val Ser Ser
 3445 3450 3455
 Ser Met Glu Phe Lys Tyr Asp Phe Asn Ser Ser Met Leu Tyr Ser Thr
 3460 3465 3470
 Ala Lys Gly Ala Val Asp His Lys Leu Ser Leu Glu Ser Leu Thr Ser
 3475 3480 3485
 Tyr Phe Ser Ile Glu Ser Ser Thr Lys Gly Asp Val Lys Gly Ser Val
 3490 3495 3500
 Leu Ser Arg Glu Tyr Ser Gly Thr Ile Ala Ser Glu Ala Asn Thr Tyr
 3505 3510 3515 3520
 Leu Asn Ser Lys Ser Thr Arg Ser Ser Val Lys Leu Gln Gly Thr Ser
 3525 3530 3535
 Lys Ile Asp Asp Ile Trp Asn Leu Glu Val Lys Glu Asn Phe Ala Gly
 3540 3545 3550
 Glu Ala Thr Leu Gln Arg Ile Tyr Ser Leu Trp Glu His Ser Thr Lys
 3555 3560 3565
 Asn His Leu Gln Leu Glu Gly Leu Phe Phe Thr Asn Gly Glu His Thr
 3570 3575 3580
 Ser Lys Ala Thr Leu Glu Leu Ser Pro Trp Gln Met Ser Ala Leu Val
 3585 3590 3595 3600
 Gln Val His Ala Ser Gln Pro Ser Ser Phe His Asp Phe Pro Asp Leu
 3605 3610 3615
 Gly Gln Glu Val Ala Leu Asn Ala Asn Thr Lys Asn Gln Lys Ile Arg
 3620 3625 3630
 Trp Lys Asn Glu Val Arg Ile His Ser Gly Ser Phe Gln Ser Gln Val
 3635 3640 3645
 Glu Leu Ser Asn Asp Gln Glu Lys Ala His Leu Asp Ile Ala Gly Ser
 3650 3655 3660
 Leu Glu Gly His Leu Arg Phe Leu Lys Asn Ile Ile Leu Pro Val Tyr
 3665 3670 3675 3680
 Asp Lys Ser Leu Trp Asp Phe Leu Lys Leu Asp Val Thr Thr Ser Ile
 3685 3690 3695
 Gly Arg Arg Gln His Leu Arg Val Ser Thr Ala Phe Val Tyr Thr Lys
 3700 3705 3710
 Asn Pro Asn Gly Tyr Ser Phe Ser Ile Pro Val Lys Val Leu Ala Asp

3715	3720	3725
Lys Phe Ile Ile Pro Gly Leu Lys Leu Asn Asp Leu Asn Ser Val Leu		
3730	3735	3740
Val Met Pro Thr Phe His Val Pro Phe Thr Asp Leu Gln Val Pro Ser		
3745	3750	3755
Cys Lys Leu Asp Phe Arg Glu Ile Gln Ile Tyr Lys Lys Leu Arg Thr		3760
3765	3770	3775
Ser Ser Phe Ala Leu Asn Leu Pro Thr Leu Pro Glu Val Lys Phe Pro		
3780	3785	3790
Glu Val Asp Val Leu Thr Lys Tyr Ser Gln Pro Glu Asp Ser Leu Ile		
3795	3800	3805
Pro Phe Phe Glu Ile Thr Val Pro Glu Ser Gln Leu Thr Val Ser Gln		
3810	3815	3820
Phe Thr Leu Pro Lys Ser Val Ser Asp Gly Ile Ala Ala Leu Asp Leu		
3825	3830	3835
Asn Ala Val Ala Asn Lys Ile Ala Asp Phe Glu Leu Pro Thr Ile Ile		3840
3845	3850	3855
Val Pro Glu Gln Thr Ile Glu Ile Pro Ser Ile Lys Phe Ser Val Pro		
3860	3865	3870
Ala Gly Ile Ala Ile Pro Ser Phe Gln Ala Leu Thr Ala Arg Phe Glu		
3875	3880	3885
Val Asp Ser Pro Val Tyr Asn Ala Thr Trp Ser Ala Ser Leu Lys Asn		
3890	3895	3900
Lys Ala Asp Tyr Val Glu Thr Val Leu Asp Ser Thr Cys Ser Ser Thr		
3905	3910	3915
Val Gln Phe Leu Glu Tyr Glu Leu Asn Val Leu Gly Thr His Lys Ile		
3925	3930	3935
Glu Asp Gly Thr Leu Ala Ser Lys Thr Lys Gly Thr Phe Ala His Arg		
3940	3945	3950
Asp Phe Ser Ala Glu Tyr Glu Glu Asp Gly Lys Tyr Glu Gly Leu Gln		
3955	3960	3965
Glu Trp Glu Gly Lys Ala His Leu Asn Ile Lys Ser Pro Ala Phe Thr		
3970	3975	3980
Asp Leu His Leu Arg Tyr Gln Lys Asp Lys Lys Gly Ile Ser Thr Ser		
3985	3990	3995
Ala Ala Ser Pro Ala Val Gly Thr Val Gly Met Asp Met Asp Glu Asp		4000
4005	4010	4015
Asp Asp Phe Ser Lys Trp Asn Phe Tyr Tyr Ser Pro Gln Ser Ser Pro		
4020	4025	4030
Asp Lys Lys Leu Thr Ile Phe Lys Thr Glu Leu Arg Val Arg Glu Ser		
4035	4040	4045
Asp Glu Glu Thr Gln Ile Lys Val Asn Trp Glu Glu Glu Ala Ala Ser		
4050	4055	4060
Gly Leu Leu Thr Ser Leu Lys Asp Asn Val Pro Lys Ala Thr Gly Val		
4065	4070	4075
Leu Tyr Asp Tyr Val Asn Lys Tyr His Trp Glu His Thr Gly Leu Thr		4080
4085	4090	4095
Leu Arg Glu Val Ser Ser Lys Leu Arg Arg Asn Leu Gln Asp His Ala		
4100	4105	4110
Glu Trp Val Tyr Gln Gly Ala Ile Arg Glu Ile Asp Asp Ile Asp Glu		
4115	4120	4125
Arg Phe Gln Lys Gly Ala Ser Gly Thr Thr Gly Thr Tyr Gln Glu Trp		
4130	4135	4140
Lys Asp Lys Ala Gln Asn Leu Tyr Gln Glu Leu Leu Thr Gln Glu Gly		
4145	4150	4155
		4160

Gln Ala Ser Phe Gln Gly Leu Lys Asp Asn Val Phe Asp Gly Leu Val
 4165 4170 4175
 Arg Val Thr Gln Glu Phe His Met Lys Val Lys His Leu Ile Asp Ser
 4180 4185 4190
 Leu Ile Asp Phe Leu Asn Phe Pro Arg Phe Gln Phe Pro Gly Lys Pro
 4195 4200 4205
 Gly Ile Tyr Thr Arg Glu Glu Leu Cys Thr Met Phe Ile Arg Glu Val
 4210 4215 4220
 Gly Thr Val Leu Ser Gln Val Tyr Ser Lys Val His Asn Gly Ser Glu
 4225 4230 4235 4240
 Ile Leu Phe Ser Tyr Phe Gln Asp Leu Val Ile Thr Leu Pro Phe Glu
 4245 4250 4255
 Leu Arg Lys His Lys Leu Ile Asp Val Ile Ser Met Tyr Arg Glu Leu
 4260 4265 4270
 Leu Lys Asp Leu Ser Lys Glu Ala Gln Glu Val Phe Lys Ala Ile Gln
 4275 4280 4285
 Ser Leu Lys Thr Thr Glu Val Leu Arg Asn Leu Gln Asp Leu Leu Gln
 4290 4295 4300
 Phe Ile Phe Gln Leu Ile Glu Asp Asn Ile Lys Gln Leu Lys Glu Met
 4305 4310 4315 4320
 Lys Phe Thr Tyr Leu Ile Asn Tyr Ile Gln Asp Glu Ile Asn Thr Ile
 4325 4330 4335
 Phe Asn Asp Tyr Ile Pro Tyr Val Phe Lys Leu Leu Lys Glu Asn Leu
 4340 4345 4350
 Cys Leu Asn Leu His Lys Phe Asn Glu Phe Ile Gln Asn Glu Leu Gln
 4355 4360 4365
 Glu Ala Ser Gln Glu Leu Gln Gln Ile His Gln Tyr Ile Met Ala Leu
 4370 4375 4380
 Arg Glu Glu Tyr Phe Asp Pro Ser Ile Val Gly Trp Thr Val Lys Tyr
 4385 4390 4395 4400
 Tyr Glu Leu Glu Glu Lys Ile Val Ser Leu Ile Lys Asn Leu Leu Val
 4405 4410 4415
 Ala Leu Lys Asp Phe His Ser Glu Tyr Ile Val Ser Ala Ser Asn Phe
 4420 4425 4430
 Thr Ser Gln Leu Ser Ser Gln Val Glu Gln Phe Leu His Arg Asn Ile
 4435 4440 4445
 Gln Glu Tyr Leu Ser Ile Leu Thr Asp Pro Asp Gly Lys Gly Lys Glu
 4450 4455 4460
 Lys Ile Ala Glu Leu Ser Ala Thr Ala Gln Glu Ile Ile Lys Ser Gln
 4465 4470 4475 4480
 Ala Ile Ala Thr Lys Lys Ile Ile Ser Asp Tyr His Gln Gln Phe Arg
 4485 4490 4495
 Tyr Lys Leu Gln Asp Phe Ser Asp Gln Leu Ser Asp Tyr Tyr Glu Lys
 4500 4505 4510
 Phe Ile Ala Glu Ser Lys Arg Leu Ile Asp Leu Ser Ile Gln Asn Tyr
 4515 4520 4525
 His Thr Phe Leu Ile Tyr Ile Thr Glu Leu Leu Lys Lys Leu Gln Ser
 4530 4535 4540
 Thr Thr Val Met Asn Pro Tyr Met Lys Leu Ala Pro Gly Glu Leu Thr
 4545 4550 4555 4560
 Ile Ile Leu

<210> 2

<211> 14070

<212> DNA

<213> Homo Sapiens

<400> 2

gaggagcccg	cccagccagc	cagggccgcg	aggccgaggg	caggccgcag	cccaggagcc	60
gccccaccgc	agctggcgat	ggacccgccg	aggcccgccg	tgctggcgct	gctggcgctg	120
cctgcgctgc	tgctgctgct	gctggcgggc	gccaggggccg	aagaggaaat	gctggaaaat	180
gtcagcctgg	tctgtccaaa	agatgcgacc	cgattcaagc	acctccggaa	gtacacatac	240
aactatgagg	ctgagagttc	cagtggagtc	cctgggactg	ctgattcaag	aagtgccacc	300
aggatcaact	gcaaggttga	gctggaggtt	ccccagctct	gcagcttcat	cctgaagacc	360
agccagtgca	tcctgaaaga	ggtgtatggc	ttcaaccctg	agggcaaagc	cttgctgaag	420
aaaaccaaga	actctgagga	gtttgctgca	gccatgtcca	ggtatgagct	caagctggcc	480
attccagaag	ggaagcaggt	tttcctttac	ccggagaaaag	atgaacctac	ttacatcctg	540
aacatcaaga	ggggcatcat	ttctgccctc	ctggttcccc	cagagacaga	agaagccaag	600
caagtgttgt	ttctggatac	cgtgtatgga	aactgctcca	ctcactttac	cgtaagacg	660
aggaagggca	atgtggcaac	agaaatatcc	actgaaagag	acctggggca	gtgtgatcgc	720
ttcaagccca	tccgcacagg	catcagccca	cttgctctca	tcaaaggcat	gacccgcccc	780
ttgtcaactc	tgtcagcag	cagccagtc	tgtcagtaca	cactggacgc	taagaggaag	840
catgtggcag	aagccatctg	caaggagcaa	cacctcttcc	tgcctttctc	ctacaagaat	900
aagtatggga	tggtagcaca	agtgcacacg	actttgaaac	ttgaagacac	accaaagatc	960
aacagccgct	tctttggtga	aggtactaag	aagatggggc	tcgcatttga	gagcaccaaa	1020
tccacatcac	ctccaaagca	ggccgaagct	gttttgaaaga	ctgtccagga	actgaaaaaa	1080
ctaaccatct	ctgagcaaaa	tatccagaga	gctaattctct	tcaataagct	ggttactgag	1140
ctgagaggcc	tcagtgatga	agcagtcaca	tctctcttgc	cacagctgat	tgagggtgct	1200
agccccatca	ctttacaagc	cttggttcag	tgtggacagc	ctcagtgtct	cactcacatc	1260
ctccagtggc	tgaacgtgt	gcatgccaac	cccccttctga	tagatgtggt	cacctacctg	1320
gtggccctga	tccccgagcc	ctcagcacag	cagctgcgag	agatcttcaa	catggcgagg	1380
gatcagcgca	gccgagccac	cttgtatgcg	ctgagccacg	cgggtcaaca	ctatcataag	1440
acaaacccta	cagggagcca	ggagctgctg	gacattgtcta	attacctgat	ggaacagatt	1500
caagatgact	gcactgggga	tgaagattac	acctatttga	ttctgcgggt	cattggaaat	1560
atgggcca	ccatggagca	gttaactcca	gaactcaagt	cttcaatcct	gaaatgtgtc	1620
caaagtacaa	agccatcact	gatgatccag	aaagctgcca	tccaggctct	gcggaatg	1680
gagcctaaag	acaaggacca	ggaggttctt	cttcagactt	tccttgatga	tgcttctccg	1740
ggagataaag	gactggctgc	ctatcttatg	ttgatgagga	gtccttcaca	ggcagatatt	1800
aacaaaattg	tccaaattct	accatgggaa	cagaatgagc	aagtgaagaa	ctttgtggct	1860
tcccatattg	ccaatatctt	gaactcagaa	gaattggata	tccaagatct	gaaaaagtta	1920
gtgaaagaag	ttctgaaaga	atctcaactt	ccaactgtca	tggacttcag	aaaattctct	1980
cggaactatc	aactctacaa	atctgtttct	attccatcac	ttgaccacgc	ctcagccaaa	2040
atagaaggga	atcttatatt	tgatccaaat	aactaccttc	ctaaagaaag	catgctgaaa	2100
actaccctca	ctgcctttgg	atttgcttca	gctgacctca	tcgagattgg	cttggaagga	2160
aaaggctttg	agccaacatt	ggaagctctt	tttggaagc	aaggattttt	cccagacagt	2220
gtcaacaaag	ctttgtactg	ggttaatggg	caagtctctg	atggtgtctc	taaggctcta	2280
gtggaccact	ttggctatac	caaagatgat	aaacatgagc	aggatatggg	aatggaata	2340
atgctcagtg	ttgagaagct	gattaaagat	ttgaaatcca	aagaagtccc	ggaagccaga	2400
gcctacctcc	gcatcttggg	agaggagctt	ggttttgcca	gtctccatga	cctccagctc	2460
ctgggaaagc	tgcttctgat	gggtgccgcg	actctgcagg	ggatcccca	gatgattgga	2520
gaggtcatca	ggaagggtc	aaagaatgac	tttttcttc	actacatctt	catggagaat	2580
gcctttgaac	tccccactgg	agctggatta	cagttgcaaa	tatcttcac	tggagtcat	2640
gtccccggag	ccaaggctgg	agtaaaactg	gaagtagcca	acatgcaggc	tgaactggtg	2700
gcaaaacct	ccgtgtctgt	ggagtgttg	acaaatatgg	gcacatcat	tccggacttc	2760
gctaggagt	gggtccagat	gaacaccaac	ttcttccacg	agtcgggtct	ggaggctcat	2820
gttgccctaa	aacctgggaa	gctgaagtgt	atcattcctt	cccaaaagag	accagtcaag	2880
ctgctcagtg	gaggcaacac	attacatttg	gtctctacca	ccaaaacgga	ggtgatccca	2940
cctctcattg	agaacaggca	gtcctggtea	gtttgcaagc	aagtctttcc	tggcctgaat	3000
tactgcacct	caggcgctta	ctccaacgcc	agctccacag	actccgcctc	ctactatccg	3060

ctgaccgggg	acaccagatt	agagctggaa	ctgaggccta	caggagagat	tgagcagtat	3120
tctgtcagcg	caacctatga	gctccagaga	gaggacagag	ccttggtgga	tacctgaag	3180
tttghtaactc	aagcagaagg	tgcgaagcag	actgaggcta	ccatgacatt	caaatataat	3240
cggcagagta	tgaccttgtc	cagtgaagtc	caaatcccg	atcttgatgt	tgacctcgga	3300
acaatcctca	gagttaatga	tgaatctact	gagggcaaaa	cgtcttacag	actcaccctg	3360
gacattcaga	acaagaaaat	tactgaggtc	gccctcatgg	gccacctaag	ttgtgacaca	3420
aaggaagaaa	gaaaaatcaa	gggtgttatt	tccatacccc	gtttgcaagc	agaagccaga	3480
agtggagatcc	tcgcccactg	gtcgccctgcc	aaactgcttc	tccaaatgga	ctcatctgct	3540
acagcttatg	gctccacagt	ttccaagagg	gtggcatggc	attatgatga	agagaagatt	3600
gaatttgaat	ggaacacagg	caccaatgta	gataccaaaa	aatgacttc	caatttccct	3660
gtggatctct	ccgattatcc	taagagcttg	catatgtatg	ctaatagact	cctggatcac	3720
agagtcctctc	aaacagacat	gactttccgg	cacgtgggtt	ccaaattaat	agttgcaatg	3780
agctcatggc	ttcagaaggc	atctgggagt	cttccttata	cccagacttt	gcaagaccac	3840
ctcaatagcc	tgaaggagtt	caacctccag	aacatgggat	tgccagactt	ccacatccca	3900
gaaaacctct	tcttaaaaag	cgatggccgg	gtcaaatata	ccttgaacaa	gaacagtttg	3960
aaaattgaga	ttcctttgcc	ttttggtggc	aaatccctcca	gagatctaaa	gatgttagag	4020
actgttagga	caccagccct	ccacttcaag	tctgtgggat	tccatctgcc	atctcgagag	4080
ttccaagtcc	ctacttttac	cattcccaag	ttgtatcaac	tgcaagtgcc	tctcctgggt	4140
gttctagacc	tctccacgaa	tgtctacagc	aacttgtaca	actggtccgc	ctcctacagt	4200
ggtggcaaca	ccagcacaga	ccatttcagc	cttcgggctc	gttaccacat	gaaggctgac	4260
tctgtggttg	acctgcttcc	ctacaatgtg	caaggatctg	gagaaacaac	atatgaccac	4320
aagaatacgt	tcacactatc	atgtgatggg	tctctacgcc	acaaatttct	agattcgaat	4380
atcaaattca	gtcatgtaga	aaaacttggg	aacaacccag	tctcaaaagg	tttactaata	4440
ttcgatgcac	ctagtctctg	gggaccacag	atgtctgctt	cagttcattt	ggactccaaa	4500
aagaaacagc	atgtgtttgt	caaagaagtc	aagattgatg	ggcagttcag	agtctcttcg	4560
ttctatgcta	aaggcacata	tggcctgtct	tgtcagaggg	atcctaacac	tggccggctc	4620
aatggagagt	ccaacctgag	gtttaactcc	tcctacctcc	aaggcaccaa	ccagataaca	4680
ggaagatatg	aagatggaac	cctctccctc	acctccacct	ctgatctgca	aagtggcatc	4740
attaaaaata	ctgcttccct	aaagtatgag	aactacgagc	tgactttaaa	atctgacacc	4800
aatgggaagt	ataagaactt	tgccacttct	aacaagatgg	atatgacctt	ctctaagcaa	4860
aatgcactgc	tgcgttctga	atatcaggct	gattacgagt	cattgaggtt	cttcagcctg	4920
ctttctggat	cactaaattc	ccatggctct	gagttaaatg	ctgacatctt	aggcactgac	4980
aaaattaata	gtggtgctca	caaggcgaca	ctaaggattg	gccaagatgg	aatatctacc	5040
agtgaacga	ccaacttgaa	gtgtagtctc	ctggtgctgg	agaatgagct	gaatgcagag	5100
cttggcctct	ctggggcatc	tatgaaatta	acaacaaatg	gccgcttcag	ggaacacaat	5160
gcaaaattca	gtctggatgg	gaaagccgcc	ctcacagagc	tatcactggg	aagtgcctat	5220
caggecatga	ttctgggtgt	cgacagcaaa	aacattttca	acttcaaggt	cagtcaagaa	5280
ggacttaagc	tctcaaatga	catgatgggc	tcatatgctg	aaatgaaatt	tgaccacaca	5340
aacagctcga	acattgcagg	cttatcactg	gacttctctt	caaaaactga	caacatttac	5400
agctctgaca	agttttataa	gcaaaactgt	aatttacagc	tacagcccta	ttctctggta	5460
actactttta	acagtgcact	gaaatacaat	gctctggatc	tcaccaacaa	tgggaaacta	5520
cggctagaac	ccctgaagct	gcatgtggct	ggtaacctaa	aaggagccta	ccaaaataat	5580
gaaataaaac	acatctatgc	catctcttct	gctgccttat	cagcaagcta	taaagcagac	5640
actgttgcta	agggttcagg	tgtggagtgt	agccatcggc	tcaacacaga	catcgctggg	5700
ctggcttcag	ccattgacat	gagcacaac	tataattcag	actcactgca	tttcagcaat	5760
gtcttccgtt	ctgtaatggc	cccgtttacc	atgaccatcg	atgcacatac	aaatggcaat	5820
gggaaactcg	ctctctgggg	agaacatact	gggcagctgt	atagcaaatt	cctgttgaaa	5880
gcgaacctc	tggcatttac	ttctctctat	gattacaaag	gctccacaag	tcatactctc	5940
gtgtctagga	aaagcatcag	tgcagctctt	gaacacaaag	tcagtgcctt	gcttactcca	6000
gctgagcaga	caggcacctg	gaaactcaag	acccaattta	acaacaatga	atacagccag	6060
gacttgatg	cttacaacac	taaagataaa	attggcgtgg	agcttactgg	acgaactctg	6120
gctgacctaa	ctctactaga	ctcccccaat	aaagtgcac	ttttactcag	tgagcccatc	6180
aatatcattg	atgcttttaga	gatgagagat	gccgttgaga	agccccaaga	atttacaatt	6240
gttgcttttg	taaagtatga	taaaaaccaa	gatgttctact	ccattaacct	cccatTTTTT	6300
gagaccttgc	aagaatatTT	tgagaggaat	cgacaaacca	ttatagttgt	actggaaaaa	6360

gtacagagaa	acctgaagca	catcaatatt	gatcaatttg	taagaaaata	cagagcagcc	6420
ctgggaaaac	ccccacagca	agctaattgat	tatctgaatt	cattcaattg	ggagagacaa	6480
gtttcacatg	ccaaggagaa	actgactgct	ctcacaaaaa	agtatagaat	tacagaaaat	6540
gatatacaaa	ttgcattaga	tgatgccaaa	atcaacttta	atgaaaaact	atctcaactg	6600
cagacatata	tgatacaatt	tgatcagtat	attaaagata	gttatgattt	acatgatttg	6660
aaaatagcta	ttgctaatat	tattgatgaa	atcattgaaa	aattaaaaag	tcttgatgag	6720
cactatcata	tccgtgtaaa	tttagtaaaa	acaatccatg	atctacattt	gtttattgaa	6780
aatattgatt	ttaacaaaag	tggaagtagt	actgcattct	ggattcaaaa	tgtggatact	6840
aagtaccaaa	tcagaatcca	gatacaagaa	aaactgcagc	agcttaagag	acacatacag	6900
aatatagaca	tccagcacct	agctggaaaag	ttaaaacaac	acattgaggc	tattgatgtt	6960
agagtgtctt	tagatcaatt	gggaactaca	atttcatttg	aaagaataaa	tgatgttctt	7020
gagcatgtca	aacactttgt	tataaatctt	attggggatt	ttgaagtagc	tgagaaaatc	7080
aatgccttca	gagccaaagt	ccatgagtta	atcgagaggt	atgaagtaga	ccaacaaatc	7140
caggttttta	tggataaatt	agtagagttg	gccaccaaat	acaagttgaa	ggagactatt	7200
cagaagctaa	gcaatgtcct	acaacaagtt	aagataaaaag	attactttga	gaaattggtt	7260
ggattttattg	atgatgctgt	caagaagcct	aatgaattat	ctttttaaac	attcattgaa	7320
gatgttaaca	aattccttga	catgttgata	aagaaattaa	agtcatttga	ttaccaccag	7380
ttgttagatg	aaaccaatga	caaaatccgt	gaggtgactc	agagactcaa	tgggtgaaat	7440
caggctcttg	aactaccaca	aaaagctgaa	gcatttaaac	tgtttttaga	ggaaaccaag	7500
gccacagttg	cagtgtatct	ggaaagccta	caggacacca	aaataacctt	aatcatcaat	7560
tggttacagg	aggctttaag	ttcagcatct	ttggctcaca	tgaaggccaa	attccgagag	7620
actctagaag	atacacgaga	ccgaatgtat	caaatggaca	ttcagcagga	acttcaacga	7680
tacctgtctc	tggtaggcca	ggtttatagc	acacttgtca	cctacatttc	tgattggttg	7740
actcttgctg	ctaagaacct	tactgacttt	gcagagcaat	attctatcca	agattgggct	7800
aaacgtatga	aagcattggg	agagcaaggg	ttcactgttc	ctgaaatcaa	gaccatcctt	7860
gggaccatgc	ctgcctttga	agtcagtctt	caggctcttc	agaaagctac	cttcagaca	7920
cctgatttta	tagtccccct	aacagatttg	aggattccat	cagttcagat	aaacttcaaa	7980
gacttaaaaa	atataaaaaat	cccatccagg	ttttccacac	cagaatttac	catccttaac	8040
accttccaca	ttccttccct	tacaattgac	tttgtagaaa	tgaaagtaaa	gatcatcaga	8100
accattgacc	agatgtgtaa	cagtgtgctg	cagtggcccg	ttccagatat	atatctcagg	8160
gatctgaagg	tggaggacat	tcctctagcg	agaatcacc	tgccagactt	ccgtttacca	8220
gaaatcgcaa	ttccagaatt	cataatccca	actctcaacc	ttaatgattt	tcaagttcct	8280
gaccttcaca	taccagaatt	ccagcttccc	cacatctcac	acacaattga	agtacctact	8340
tttggcaagc	tatacagtat	tctgaaaaatc	caatctcctc	ttttcacatt	agatgcaaat	8400
gctgacatag	ggaatggaac	cacctcagca	aacgaagcag	gtatcgagc	ttccatcact	8460
gccaaaggag	agtcctaaat	agaagttctc	aatttttgatt	ttcaagcaaa	tgcacaactc	8520
tcaaacctta	agattaatcc	gctggctctg	aaggagtcag	tgaagttctc	cagcaagtac	8580
ctgagaacgg	agcatgggag	tgaaatgctg	ttttttggaa	atgctattga	gggaaaatca	8640
aacacagtgg	caagtttaca	cacagaaaaa	aatacactgg	agcttagtaa	tggagtgtat	8700
gtcaagataa	acaatcagct	taccttggt	agcaacacta	aatacttcca	caaattgaac	8760
atccccaaac	tggacttctc	tagtcaggct	gacctgcgca	acgagatcaa	gacactggtg	8820
aaagctggcc	acatagcatg	gacttcttct	ggaaaagggt	catggaaatg	ggcctgcccc	8880
agattctcag	atgagggaac	acatgaatca	caaattagtt	tcaccataga	aggacccttc	8940
acttcccttg	gactgtccaa	taagatcaat	agcaaacacc	taagagtaaa	ccaaaacttg	9000
gtttatgaat	ctggctccct	caacttttct	aaacttgaaa	ttcaatcaca	agtcgattcc	9060
cagcatgtgg	gccacagtgt	tctaactgct	aaaggcatgg	cactgttttg	agaagggaag	9120
gcagagttta	ctgggaggca	tgatgtctat	ttaaatggaa	aggttatttg	aactttgaaa	9180
aattctcttt	tcttttcagc	ccagccattt	gagatcacgg	catccacaaa	caatgaaggg	9240
aatttgaaaag	tctgttttcc	attaaggtta	acagggaaga	tagacttcct	gaataactat	9300
gcactgtttc	tgagtcccag	tgcccagcaa	gcaagttggc	aagtaagtgc	taggttcaat	9360
cagtataagt	acaacaaaaa	tttctctgct	ggaaacaacg	agaacattat	ggaggccccat	9420
gtaggaaata	atggagaagc	aaatctggat	tctttaaaca	ttcctttaac	aattcctgtg	9480
atgcgtctac	cttacacaat	aatcacaaat	cctccactga	aagatttctc	tctatgggaa	9540
aaaacaggct	tgaagggaatt	cttgaaaacg	acaaagcaat	catttgattt	aagtgtaaaa	9600
gctcagtata	agaaaaacaa	acacaggcat	tccatcacia	atcctttggc	tgtgctttgt	9660

gagtttatca	gtcagagcat	caaatccttt	gacaggcatt	ttgaaaaaaa	cagaaacaat	9720
gcattagatt	ttgtcaccaa	atcctataat	gaaacaaaaa	ttaagtttga	taagtacaaa	9780
gctgaaaaat	ctcacgacga	gctccccagg	acctttcaaa	ttcctggata	cactgttcca	9840
gttgtcaatg	ttgaagtgtc	tccattcacc	atagagatgt	cggcattcgg	ctatgtgttc	9900
ccaaaagcag	tcagcatgcc	tagtttctcc	atcctagggt	ctgacgtccg	tgtgccttca	9960
tacacattaa	tccatgccatc	attagagctg	ccagtccttc	atgtccctag	aaatctcaag	10020
ctttctcttc	cagattttcaa	ggaattgtgt	accataagcc	atatttttat	tcttgccatg	10080
ggcaatatta	cctatgattt	ctcctttaaa	tcaagtgtca	tcacactgaa	taccaatgct	10140
gaacttttta	accagtcaga	tattgttgct	catctccttt	cttcattcttc	atctgtcatt	10200
gatgcactgc	agtacaaatt	agagggcacc	acaagattga	caagaaaaag	gggattgaag	10260
ttagccacag	ctctgtctct	gagcaacaaa	tttgtggagg	gtagtataaa	cagtactgtg	10320
agcttaacca	cgaataatat	ggaagtgtca	gtggcaacaa	ccacaaaagc	ccaaattcca	10380
atthtgagaa	tgaattttcaa	gcaagaactt	aatggaaata	ccaagtcaaa	acctactgtc	10440
tcttctctca	tggaaatttaa	gtatgatttc	aattcttcaa	tgtcttactc	taccgctaaa	10500
ggagcagttg	accacaagct	tagcttggaa	agcctcacct	cttacttttc	cattgagtca	10560
tctaccaaag	gagatgtcaa	gggttcgggt	ctttctcggg	aatattcagg	aactattgct	10620
agtgaaggca	acacttactt	gaattccaag	gtcacacggg	cttcagtcaa	gctgcagggc	10680
acttccaaaa	ttgatgatat	ctggaacctt	gaagtaaaaag	aaaattttgc	tggagaagcc	10740
acactccaac	gcataatatt	cctctgggag	cacagtacga	aaaaccactt	acagctagag	10800
ggcctctttt	tcaccaacgg	agaacataca	agcaaaagca	ccctggaact	ctctccatgg	10860
caaatgtcag	ctcttgttca	ggtccatgca	agtcagccca	gttccttcca	tgatttccct	10920
gaccttggcc	aggaagtggc	cctgaatgct	aacactaaga	accagaagat	cagatggaaa	10980
aatgaagtc	ggattcatte	tgggtctttc	cagagccagg	tgcagctttc	caatgaccaa	11040
gaaaaggcac	accttgacat	tgcaggatcc	ttagaaggac	acctaagggt	cctcaaaaat	11100
atcatcctac	cagtcctatga	caagagctta	tgggattttc	taaagctgga	tgtaaccacc	11160
agcattggta	ggagacagca	tctctgtgtt	tcaactgcct	ttgtgtacac	caaaaacccc	11220
aatggctatt	catttctccat	ccctgtaaaa	gttttggctg	ataaattcat	tattcctggg	11280
ctgaaactaa	atgatctaaa	ttcagttctt	gtcatgccta	cgttccatgt	cccatttaca	11340
gatcttcagg	ttccatcgtg	caaacttgac	ttcagagaaa	tacaaatcta	taagaagctg	11400
agaacttcat	catttgcctt	caacctacca	acactccccg	aggtaaaaat	ccctgaagtt	11460
gatgtgttaa	caaaatattc	tcaaccagaa	gactccttga	ttcccttttt	tgagataaac	11520
gtgcctgaat	ctcagttaac	tgtgtcccag	ttcacgcttc	caaaaagtgt	ttcagatggc	11580
attgtctgct	tggatctaaa	tgcagttagc	aacaagatcg	cagactttga	gttgccacc	11640
atcatctgtc	ctgagcagac	cattgagatt	ccctccatta	agttctctgt	acctgctgga	11700
attgccattc	cttcttttca	agcactgact	gcacgctttg	aggtagactc	tccctgttat	11760
aatgccactt	ggagtgccag	tttgaaaaac	aaagcagatt	atgttgaaac	agtcctggat	11820
tccacatgca	gctcaaccgt	acagtcccta	gaatatgaac	ttaatgtttt	gggaacacac	11880
aaaatcgaa	atggtacgtt	agcctctaag	actaaaggaa	catttgcaca	ccgtgacttc	11940
agtgcagaat	atgaagaaga	tggcaaatat	gaaggacttc	aggaatggga	aggaaaagcg	12000
cacctcaata	tcaaaaagccc	agcgttcacc	gatctccatc	tgcgctacca	gaaagacaag	12060
aaaggcatct	ccacctcagc	agcctcccca	gccgtaggca	ccgtgggcat	ggatatggat	12120
gaagatgacg	actttttctaa	atggaacttc	tactacagcc	ctcagtcctc	tccagataaa	12180
aaactcacca	tattcaaaac	tgagttgagg	gtccgggaat	ctgatgagga	aactcagatc	12240
aaagttaatt	gggaagaaga	ggcagcttct	ggcttgctaa	cctctctgaa	agacaacgtg	12300
cccaaggcca	caggggtcct	ttatgattat	gtcaacaagt	accactggga	acacacaggg	12360
ctcacctga	gagaagtgtc	ttcaaaagctg	agaagaaatc	tgcaggacca	tgtgtagtgg	12420
gtttatcaag	gggccattag	ggaaattgat	gatatcgacg	agaggttcca	gaaaggagcc	12480
agtgggacca	ctgggacctt	ccaagagtgg	aaggacaagg	cccagaatct	gtaccaggaa	12540
ctgttgactc	aggaaggcca	agccagtttc	cagggactca	aggataacgt	gtttgatggc	12600
ttgggtacag	ttactcaaga	attccatag	aaagtcaagc	atctgattga	ctcactcatt	12660
gattttctga	acttccccag	attccagttt	ccggggaaac	ctgggatata	cactagggag	12720
gaactttgca	ctatgttcat	aagggaggta	gggacggtac	tgtcccaggt	atattcgaaa	12780
gtccataatg	gttcagaaat	actgttttcc	tatttccaag	acctagtgat	tacacttcct	12840
ttcgagttaa	ggaaacataa	actaatagat	gtaattctcg	tgtataggga	actgttgaaa	12900
gatttatcaa	aagaagccca	agaggtattt	aaagccattc	agtctctcaa	gaccacagag	12960

```

gtgctacgta atcttcagga ccttttacaa ttcattttcc aactaataga agataacatt 13020
aaacagctga aagagatgaa atttacttat cttattaatt atatccaaga tgagatcaac 13080
acaatcttca atgattatat cccatatgtt tttaaattgt tgaaagaaaa cctatgcctt 13140
aatcttcata agttcaatga atttattcaa aacgagcttc aggaagcttc tcaagagtta 13200
cagcagatcc atcaatacat tatggccctt cgtgaagaat attttgatcc aagtatagtt 13260
ggctggacag tgaatatatta tgaacttgaa gaaaagatag tcagtctgat caagaacctg 13320
ttagtgtctc ttaaggactt ccattctgaa tatattgtca gtgcctctaa ctttacttcc 13380
caactctcaa gtcaagttga gcaatttctg cacagaaata ttcaggaata tcttagcatc 13440
cttaccgatc cagatggaaa agggaaagag aagattgcag agctttctgc cactgctcag 13500
gaaataatta aaagccaggc cattgcgacg aagaaaataa tttctgatta ccaccagcag 13560
tttagatata aactgcaaga tttttcagac caactctctg attactatga aaaatttatt 13620
gctgaatcca aaagattgat tgacctgtcc attcaaaact accacacatt tctgatatac 13680
atcacggagt tactgaaaaa gctgcaatca accacagtca tgaaccctca catgaagctt 13740
gctccaggag aacttactat catcctctaa ttttttaaaa gaaatcttca tttattcttc 13800
ttttccaatt gaactttcac atagcacaga aaaaattcaa aatgcctata ttgatcaaac 13860
catacagtga gccagccttg cagtaggcag tagactataa gcagaagcac atatgaactg 13920
gacctgcacc aaagctggca ccagggtctg gaagggtctc gaactcagaa ggatggcatt 13980
ttttgcaagt taaagaaaat caggatctga gttattttgc taaacttggg ggaggaggaa 14040
caaataaatg gagtctttat tgtgtatcat 14070

```

```

<210> 3
<211> 3805
<212> DNA
<213> Homo Sapiens

```

```

<220>
<221> exon
<222> (71)...(114)
<223> Exon 1

<221> exon
<222> (872)...(937)
<223> Exon 2

<221> exon
<222> (2031)...(2223)
<223> Exon 3

```

```

<221> exon
<222> (2805)...(3663)
<223> Exon 4

```

```

<400> 3
cctatccctg ggggaggggg cgggacaggg ggagccctat aattggacaa gtctggggtc 60
cttgagtcct actcagccc agcggagggtg aaggacgtcc tccccagga gccggtgaga 120
agcgcagtcg ggggcacggg gatgagctca ggggcctcta gaaagagctg ggaccctggg 180
aagccctggc ctccaggtag tctcaggaga gctactcggg gtcgggcttg gggagaggag 240
gagcgggggt gaggcaagca gcaggggact ggacctggga agggctgggc agcagagacg 300
acccgaccgg ctagaagggtg ggggtggggag agcagctgga ctgggatgta agccatagca 360
ggactccacg agttgtcact atcattatcg agcacctact ggggtgtcccc agtgtcctca 420
gatctccata actggggagc caggggcagc gacacggtag ctagccgtcg attggagaac 480
tttaaaatga ggactgaatt agctcataaa tggaaacagg cgcttaactg tgaggttga 540
gcttagaatg tgaagggaga atgaggaatg cgagactggg actgagatgg aaccggcggt 600

```

ggggaggggg	tggggggatg	gaatttgaac	ccggggagag	gaagatggaa	ttttctatgg	660
agggcgacct	ggggatgggg	agataagaga	agaccaggag	ggagttaa	agggaaatggg	720
ttggggcgcg	cttggtaaat	gtgctgggat	taggctgttg	cagataatgc	aacaaggctt	780
ggaaggctaa	cctggggtga	ggccgggttg	ggggcgctgg	gggtgggagg	agtcctcact	840
ggcggttgat	tgacagtttc	tccttcccca	gactggccaa	tcacaggcag	gaagatgaag	900
gttctgtggg	ctgcgttgct	ggtcacattc	ctggcaggta	tgggggcggg	gcttgctcgg	960
ttcccccgcg	tcctccccct	ctcatcctca	cctcaacctc	ctggcccat	tcagacagac	1020
cctgggcccc	ctcttctgag	gcttctgtgc	tgcttccctg	ctctgaacag	cgatttgacg	1080
ctctctgggc	ctcggtttcc	cccatccttg	agataggagt	tagaagttgt	tttgttggtg	1140
ttgtttgttg	ttgttggttt	gtttttttga	gatgaagtct	cgctctgtcg	cccaggctgg	1200
agtgcagtgg	cgggatctcg	gctcactgca	agctccgcct	cccaggcca	cgccattctc	1260
ctgcctcagc	ctcccaagta	gctgggacta	caggcacatg	ccaccacacc	cgactaactt	1320
ttttgtattt	tcagtagaga	cggggtttca	ccatgttggc	caggctggtc	tggaaactcct	1380
gacctcaggt	gatctgcccc	tttcgatctc	ccaaagtgtc	gggattacag	gcgtgagcca	1440
ccgcacctgg	ctgggagtta	gaggtttcta	atgcattgca	ggcagatagt	gaataaccaga	1500
cacggggcag	ctgtgatctt	tattctccat	cacccccaca	cagcctgtgc	tggggcacac	1560
aaggacactc	aatacatgct	tttccgctgg	gccggtggct	cacccctgta	atcccagcac	1620
tttgggaggg	caaggtggga	ggatcacttg	agcccaggag	ttcaacacca	gcctgggcaa	1680
catagtga	ccctgtctct	actaaaaata	caaaaattag	ccaggcatgg	tgccacacac	1740
ctgtgctctc	agctactcag	gaggctgagg	caggaggatc	gcttgagccc	agaaggca	1800
ggttgcagtg	aacctgttcc	aggccgctgc	actccagcct	gggtgacaga	gcaagacctt	1860
gtttataaat	acataatgct	ttccaagtga	ttaaaccgac	ttccccctca	ccctgcccac	1920
catggctcca	aagaagcatt	tgtggagcac	cttctgtgtg	cccctaggta	gctagatgcc	1980
tggacggggg	cagaaggacc	ctgacccgac	cttgaacttg	ttccacacag	gatgccaggc	2040
caaggtggag	caagcggtgg	agacagagcc	ggagcccag	ctgcgccagc	agaccgagtg	2100
gcagagcggc	cagcgctggg	aactggcact	gggtcgcttt	tgggattacc	tgcgctgggt	2160
gcagacactg	tctgagcagg	tgcaggagga	gctgctcagc	ttccaggcca	cccagggaact	2220
gaggtgagtg	ttcccatcct	ggcccttgac	cctcctgggt	ggcggtata	cctccccagg	2280
tccagggtttc	attctgcccc	tgtcgctaag	tcttgggggg	cctgggtctc	tgctggttct	2340
agcttcctct	ttccatttct	gactcctggc	tttagctctc	tggaaattctc	ttctctcagct	2400
ttgtctctct	ctcttccctt	ctgactcagt	ctctcacact	cgctcctggct	ctgtctctgt	2460
ccttccctag	ctcttttata	tagagacaga	gagatggggg	ctcactgtgt	tgcccaggct	2520
ggtcttgaac	ttctgggctc	aagcgatcct	ccgcctcgg	cctcccaaag	tgctgggatt	2580
agaggcatga	gcaccttgcc	cggcctccta	gctccttctt	cgctctctgcc	ttctgcccct	2640
gcactctgtc	tctgcatctg	tctctgtctc	cttctctcgg	cctctgcccc	gttctctctc	2700
ttccctcttg	gtctctctgg	ctcatcccca	tctcgccgcg	cccattcccag	cccttctccc	2760
ccgcctcccc	actgtgcgac	accctcccgc	cctctcggcc	gcaggggcgt	gatggacgag	2820
accatgaagg	agttgaaggc	ctacaaatcg	gaactggagg	aacaactgac	cccgggtggcg	2880
gaggagacgc	gggcacggct	gtccaaggag	ctgcaggcgg	cgcaggcccg	gctggggcgcg	2940
gacatggagg	acgtgtgcgg	ccgcctgggt	cagtaccgcg	gcgagggtgca	ggccatgctc	3000
ggccagagca	ccgaggagct	gcgggtgcgc	ctcgccctcc	acctgcgcaa	gctgcgtaag	3060
cggctcctcc	gcgatgccga	tgacctgcag	aagcgctgg	cagtgtacca	ggccggggcc	3120
cgcgaggggc	ccgagcgcg	cctcagcgcc	atccgcgagc	gcctggggcc	cctgggtggaa	3180
caggggccgcg	tgcggggccg	cactgtgggc	ttcctggccg	gccagccgct	acaggagcgg	3240
gcccaggcct	ggggcgagcg	gctgcgcgcg	cggatggagg	agatgggag	ccggaccgcc	3300
gaccgcctgg	acgaggtgaa	ggagcagggt	gcggagggtg	gcgccaagct	ggaggagcag	3360
gcccagcaga	tacgcctgca	ggccgaggcc	ttccaggccc	gcctcaagag	ctgggttcgag	3420
cccttgggtg	aagacatgca	gcgccagtgg	gccgggctgg	tggagaaggt	gcagggtgcc	3480
gtgggcacca	gcgcgcggcc	tgtgcccagc	gacaaatcact	gaacgcgcaa	gctgcgagcc	3540
atgcgacccc	acgccacccc	gtgcctcctg	cctccgcgca	gcctgcagcg	ggagaccctg	3600
ttcccgcccc	agccgtcctc	ctgggggtgga	ccctagttta	ataaagattc	accaagtttc	3660
acgcactctgc	tggcctcccc	ctgtgatttc	ctctaagccc	cagcctcagt	ttctctttct	3720
gcccacatac	tgccacacaa	ttctcagccc	cctcctctcc	atctgtgtct	gtgtgtatct	3780
ttctctctgc	cctttttttt	ttttt				3805

18

<210> 4
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 4
agtctggatg ggtaagccgc cctca 25

<210> 5
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 5
ctgggctggc ttaagccatt gacat 25

<210> 6
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 6
gctctctggg gataacatac tgggc 25

<210> 7
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 7
gatgccgttg agtagcccca agaat 25

<210> 8
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 8
gagaggaatc gataaaccat tatag 25

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 9
tgtaagaaaa taaagagcag ccctg 25

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 10
gcagccctgg gataactccc acagc 25

<210> 11
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 11
gcaagctaatt gattagctga attcattcaa t 31

<210> 12
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 12
caagtttcac atgcctagga gaaactgact g 31

<210> 13
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 13
atatacaaat tgcattgagat gatgccaaaa t 31

<210> 14
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 14
aaactatctc aactgtagac atatatgata c 31

<210> 15
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 15
gctaataatta ttgattaaat cattgaaatt a 31

<210> 16
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 16
tgatgagcac tagcatatcc gtgta 25

<210> 17
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 17
ctgcagcagc tttagagaca catatc 25

<210> 18
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 18
aacagtgagc tgtagtgccc cgttc 25

21

<210> 19
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 19
cagacttccg ttaaccagaa atcgc 25

<210> 20
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 20
aaagggcat ggtaatgggc ctgcc 25

<210> 21
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 21
acatatatga tataatttga tcagt 25

<210> 22
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 22
atggaggacg tgtgcggccg cctgg 25

<210> 23
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 23
gacctgcaga agtgcctggc agtgt 25

22

<210> 24
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment

<400> 24
 gacctgcaga agcgctggc agtgt 25

<210> 25
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment

<400> 25
 taaggtcagg agtttgagac cagcc 25

<210> 26
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment-DNA and RNA

<400> 26
 agucuggaug ggtaagccgc ccuca 25

<210> 27
 <211> 39
 <212> DNA
 <213> R. Rattus

<400> 27
 aaagattcat gtgaaggaga tagtggggga ccccatgtt 39

<210> 28
 <211> 13
 <212> PRT
 <213> R. Rattus

<400> 28
 Lys Asp Ser Cys Glu Gly Asp Ser Gly Gly Pro His Val
 1 5 10

<210> 29
 <211> 39
 <212> DNA
 <213> R. Rattus

<400> 29
aaagattcat gtgaaggaga tcgtggggga ccccatgtt 39

<210> 30
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 30
gggggtcccc acgatctcct tcacattttu gugaaggaga tcgtggggga ccccgcggt 60
tttcgcg 68

<210> 31
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 31
tgtgaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu cacagcggt 60
tttcgcg 68

<210> 32
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 32
tgtgaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu cacagcggt 60
tttcgcg 68

<210> 33
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 33
tgtcaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu gacagcggt 60
tttcgcg 68

<210> 34
<211> 25
<212> DNA
<213> Artificial Sequence

24

<220>
<223> PCR Primer

<400> 34
cattgctgac aaggaataca cgaac 25

<210> 35
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 35
atttgcccttt cattgcacac tcttc 25

<210> 36
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 36
attgccttgc tggaactgga taac 24

<210> 37
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 37
ttgcctttca ttgcacattc ttcac 25

<210> 38
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 38
aaggagatag tggggga 17

<210> 39
<211> 17
<212> DNA
<213> Artificial Sequence

25

<220>
<223> Oligonucleotide Probe

<400> 39
aaggagatcg tggggga 17

<210> 40
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 40
aaggagatag tggggga 17

<210> 41
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 41
aaggagatcg tggggga 17

<210> 42
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 42
attgccttgc tgggaactgga taaac 25

<210> 43
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 43
ttgcctttca ttgcacattc ttcac 25

<210> 44
<211> 17
<212> DNA
<213> Artificial Sequence

26

<220>
 <223> Oligonucleotide Probe

 <400> 44
 aaggagatag tggggga 17

 <210> 45
 <211> 17
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Oligonucleotide Probe

 <400> 45
 aaggagatcg tggggga 17

 <210> 46
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer

 <400> 46
 gttgaccgag ccacatgcct tag 23

 <210> 47
 <211> 68
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Therapeutic Oligonucleotide-DNA and RNA

 <400> 47
 ctgggctggc ttaagccatt gacatuuuua ugucaauggc uuaagccagc ccaggcgcggu 60
 uuucgcgc 68

 <210> 48
 <211> 88
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Therapeutic Oligonucleotide-DNA and RNA

 <400> 48
 gacaagtttc acatgcctag gagaaactga ctgctuuaa gcagucaguu ucucctaggc 60
 augugaaacu ugucgcgcgu uuucgcgc 88

 <210> 49
 <211> 76
 <212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 49

ccctgaatgt cctggaaatg actgccgatt ttttaucttca gucattttcca ggacauucag 60
gggcgcgttt tcgcgc 76

<210> 50

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 50

taagtttctt aactgggatt attagctggg gttttcccca gctaataatc ccagttaaga 60
aacuuagcgc gttttcgcgc 80

<210> 51

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 51

gtttcttaac tgggattatt agctgttttc agcuaataat ccaguuuag aaacgcgcgt 60
tttcgcgc 68

<210> 52

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 52

taagtttctt aactgggatt attagctggg guuuucccca gcuaauaaucc ccaguuuaga 60
aacuuagcgc guuuucgcgc 80

<210> 53

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 53

28

taagtttctt aactgggatt attagctggg guuuucccca gcuaauaauc ucaguuaaga 60
aacuuagcgc guuuucgcgc 80

<210> 54
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 54
ggcgaggaca tgtgcgaccg ggcgc 25

<210> 55
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 55
gagatgcgcg actgcgcgcg cgccc 25

<210> 56
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 56
cgcgaccgcg cgtgcgcgca tgtgg 25

<210> 57
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 57
agcgaccagc tgtgccagcg cttgg 25

<210> 58
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 58
gagctgcgcc agtgcttgcc cgcgc

25

<210> 59
<211> 8966
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (703)...(748)

<221> CDS
<222> (1678)...(2280)

<400> 59
ctaaagaaga gcactgggtg gagacacagg cgggggaagg gggaggggag tgaagtagtc 60
tccttggaaat gctgggtggt ggggaggcag tctccttggg ggaggagtcc cagcgtccct 120
cccctcccct cctctgccaa cacaatggac aatggcaact gccacacac tcccatggag 180
gggaagggga tgagtgcagg gaaccccgac cccacccggg agacctgcaa gcctgcagca 240
ctcccctccc gcccccactg aacctttgac ccttgccctg cacgccccgc agcttgctgt 300
ttgcccactc ctatttgccc agtcccagg acagagctga tccctgaact ctttaagtcc 360
acattgccag gaccagttag cagcaacagg gccagggtcg ggcttatcag cctcccagcc 420
cagacctggg ctgcagacat aaataggccc tgcaagagct ggctgcttag agactgcgag 480
aaggaggtgc gtccctgctgc ctgccccggc actctggctc cccagctcaa gggttcaggcc 540
ttgccccagg cggggcctct ggggtacctga ggtcttctcc cgtctgtgct ccttctcctc 600
acctggctgc aatgagtggg ggagcacggg gcttctgcat gctgaaggca cccactcag 660
ccaggccctt cttctcctcc aggtccccca cggcccttca ggatgaaagc tgcggtgctg 720
accttgcccg tgctcttctc gacgggtagg tgtcccctaa cctaggagcc aaccatcggg 780
gggctttctc cctaaatccc cgtggccccc cctcctgggc agaggcagca ggtttctcac 840
tgccccctc tccccaccc ccaagcttgg cctttcggct cagatctcag cccacagctg 900
gcctgatctg ggtctcccct cccaccccca gggagccagg ctggcattt ctggcagcaa 960
gatgaacccc cccagagccc ctggggtcga gtgaaggacc tggccactgt gtacgtggat 1020
gtgctcaaag acagcggcag agactatgtg tccagtttg aaggctccgc cttgggaaaa 1080
cagctaaagt aaggaccag cctggggttg agggcagggg cagggggcag aggcctgtgg 1140
gatgatgttg aagccagact ggccagctcc tcacctaata tctgatgagc tgggccccac 1200
agatggtctg gatggagaaa ccggaatgga tctccaggca gggtcacagc ccatgtcccc 1260
tgcaaaggac agaccagggc tgcccagatg gtgatcacag agccacattg tgcttgcaag 1320
tgtagcaagc ccttttccct tcttcaccac ctctctgct cctgcccagc aagactgtgg 1380
gctgtcttcg gagaggagaa tgcgctggag gcatagaagc gaggtccttc aagggccac 1440
tttgagagac aacgtaactg ggcaccagtc ccagctctgt ctccctttta gctcctctct 1500
gtgcctcggc ccagctgcac aacggggcat ggcttgccgg ggcaggggtg ttggttgaga 1560
gtgtactgga aatgctaggc cactgcacct ccgaggacag gtgtcaccca gggctcacc 1620
ctgataggct ggggcgctgg gagggcagcc ctcaaccctt ctgtctcacc ctccagccta 1680
aagctccttg acaactggga cagcgtgacc tccaccttca gcaagctgag cgaacagctc 1740
ggcctgtgta cccaggagtt ctgggataac ctggaaaagg agacagaggg cctgaggcaa 1800
gagatgagca aggatctgga ggagggtgaag gccaaagtgc agccctacct ggacgacttc 1860
cagaagaagt ggcaggagga gatggagctc taccgccaga aggtggagcc gctgcgcgca 1920
gagctccaag agggcgcgcg ccagaagctg cagcagctgc aagagaagct gagccactg 1980
ggcgaggaga tgcgcgaccg cgcgcgcgcc catgtggacg cgtgcgcgac gcatctggcc 2040
ccctacagcg acgagctgag ccagcgttgg gccgcgcgcc ttgaggctct caaggagaac 2100
ggcgcgccca gactggccga gtaccacgcc aaggccaccg agcatctgag cacgctcagc 2160
gagaaggcca agcccgcgct cgaggacctc cgccaaggcc tgcctgccgt gctggagagc 2220
ttcaaggctc gcttctctgag cgtctctgag gagtacacta agaagctcaa caccagtgta 2280
ggcgccccgc gccgcccccc tccccgggtg tcagaataaa cgtttccaaa gtgggaagca 2340

gcttctttct	tttgggagaa	tagagggggg	tgcgggggaca	tccggggggag	cccgggaggg	2400
gcctttggcc	ctggagcagg	gacttctctgc	cggatctcaa	caactccgtg	cccagactgg	2460
acgtcttagg	gccaaagatcg	acgttggagg	acctgtgga	cgcntggctg	cttacgagtg	2520
agggagtaga	gtctgcctta	gcaaggctca	agtagaaagg	aagtcacagc	ggacnaggca	2580
aagccacaga	caatccaagg	ccaggtgccc	tgaagggggc	tcaaacaagg	cctgcagcct	2640
gtctgaggcg	ggccaggaaa	cagcggttgc	tttagctggg	agcagtcggg	ttccccgtcc	2700
ccagaggtgt	gtccgtatag	agccttctcc	agcccagccg	ctgtcagcgg	gcgggacgga	2760
gcggggcgcc	tcaggggagcc	agccactggg	attgggggtt	ggtcccgggt	gcaagtgaag	2820
cgcttggagt	ttgcgcctgt	cctcctttac	taattcaaaa	acctctcaa	cagacacttc	2880
ccttttcttc	tcacaaggcc	agtatcccc	tcccactact	cccatcccg	ccagaaacag	2940
ccgcggcttc	ctcaggcaca	gcagtggaag	ccagtcctcc	acccctcg	gctccatgcc	3000
atgccacccc	ctctttctgc	cagccctggc	agaagctggc	ctgagtaaga	aaattcacca	3060
ccacctcttg	caggtagcatt	tttatttcca	agatgctctc	atatctgtgc	tctcactgca	3120
tcttcccttc	cccacatcct	ggctagattg	ccatcagacg	cagagcatgg	atgaggacac	3180
tgaagcctgg	acctgtgacg	tcgcttggcc	agtgaacagc	aggatgggct	aggcagcgct	3240
tttttagacc	tgcaccctcg	gccatccatg	attattgaaa	agagtgtg	ggtcgggtgc	3300
ggtggtctca	gcctgtaatc	ccagcacttt	gggaggctga	ggtgggctga	tcacttcagg	3360
gccaggagtt	tgagaccagc	ctggccaata	tggtgaaacc	ctgtctctac	taaaaataca	3420
aaaaaaaaat	cagctggcat	ggtggcttgc	acccgtaatc	ccagctacta	ggaaggctga	3480
ggcaggagaa	tcgcttgaa	ctgggaggca	gaggtcacag	tgagccgaaa	tcatgccact	3540
gcactccagc	ctgggagcag	gagcaagact	ccagctaaaa	aaaaaaaaaa	aaaaaaaaag	3600
agtgtgtggc	ctggcactca	agttcacatg	ggtgtgcagg	catgcctgtg	tattctcaca	3660
tgacctccct	gctcacggtc	cctccttgca	ctcatgtctg	aatgtccccg	ctgcacgcac	3720
atggcttcac	agatctggca	gtgccttccc	tacctctct	ctgcagggcc	ttttgcccc	3780
tcattgaggc	ccctggataa	tcggctccat	ccccatgtcc	ccatctccag	tgtatcttag	3840
ctacctaggt	taaaggagtg	ggcttttttag	ttcctaacct	tccagagcta	caacagcagt	3900
catccagcca	ggtctgggtg	ggaacatttt	ctagatcagg	gtgctgagat	ctctcagccc	3960
agagagaagc	cctggggaat	tttcagagag	aaagcagctc	ccagggtggg	ctggatgtac	4020
tgatgccact	gagatctgta	aaggagtccc	taacacctga	cataggagtg	acaaaactgt	4080
tttctgcacc	aactgagcag	aatacacgca	gctgacctgg	gctcaaggct	tggccctgcc	4140
acgtgctggc	tctgtgatgc	tggccaagtg	ccttcgcctc	tccggggccac	agttttttga	4200
tctgaagagt	ggagccctac	tcaagccatc	gggctctctg	gggctctctg	acctgacatc	4260
tttcgggtgg	tggggacaca	aaggaaagcag	cctctattgg	gagaccttgt	gcttcttttt	4320
ggtccaggga	caactgcccc	caccactcca	gtccggtccc	aaggggccag	tcagctcaac	4380
tgtaatcatg	acaacattga	tcaagcatct	ttacgtgcag	gtgctgtgcc	aaacggttcg	4440
aacgctctct	catttcaatc	tcacggcaaa	cctacgggtg	aggggggtacg	gttgtatcca	4500
ctttacatgt	aagaaactga	ggctgatatc	aagtgggtga	gccaagaata	gtgcctcgct	4560
gcatcttact	ccaacctcta	gcccattccg	cctcctccct	tcacgtgcgc	ctaagagggc	4620
taggtggcct	ggatagggga	ggtcagctcc	acagttttga	gtaaacacac	acagtcctca	4680
ctctgatgac	acttaagtgc	caggcatagt	ggctggcatg	gggcacacac	tcaagtcatg	4740
ttgtgcagca	cctaacagtt	tatcaaagta	tcagcaaaact	tattgtcctg	tttgaccttc	4800
cgcacaaaagc	tgtcaaggaa	ggcagggtac	ggagggtgat	tcctacctta	gagatgaaga	4860
aactgaggcc	cagagactag	cccagctacc	agaagggtga	tagagcgctg	gcctccatgc	4920
ctgcctgacc	tggagtctgt	ccagtgccca	cccacagaac	agcctcggcc	ctttcccatg	4980
cccagacaca	gatggcacac	ttgcgacggc	ccactcatag	cagcttcttg	tccagcttta	5040
ttgggaggcc	agcatgacctg	gagggggggc	aggcatgagg	tggggtagga	gagcactgag	5100
aatactgtcc	cttttaagca	acctacaggg	gaagccctgg	agattgcagg	acccaaggag	5160
ctggcaggat	ggataggcag	gtggacttgg	ggtattgagg	tctcaggcag	ccacggctga	5220
agttgggtctg	acctcagggt	ccaaatccca	gaactcagag	aacttgcct	taacgggtgct	5280
ccagtagtct	ttcagggaa	tgaagccatc	ggtcaccag	cccctaaatc	agtcagggga	5340
agcaacagag	cagggcatga	gaactcctct	gtaggcaacc	atgggaacca	cacccatgtc	5400
cccactggac	gacaccagtc	aggaccacac	caccctctca	acttcaactg	acgacagccc	5460
tgagacctca	ggcaggaatc	ccccacagct	gccaccctgg	gagaagagat	atccttgacg	5520
gaacccccagc	acaagtcaaa	ccctgccatc	tccaggtcac	ctcagatgtt	tatgcccctg	5580
ggcctgaggc	acaaagtgc	aggggtggga	gatgtgaaag	gtcaaggctg	tcattgtttt	5640

ctgtgcaaac	agcaccacct	ggagttgcac	aacctggtgg	ctctgagcag	ggtaggacag	5700
agggaggcag	cctctcattt	ggaaagtcac	tggaggattg	attagttgtg	tgatctgggg	5760
caggtcacct	aatcgctctg	agcctcaatt	ttctcatctg	caaagtgaga	aaataacacc	5820
taccccaaac	ccggttcttg	ggactaagaa	tgtttatgaa	cacctctgct	atgccagcta	5880
atgccagtca	gggtgagggtg	gagaagggag	taggggagag	gagagtactg	atgggcaggg	5940
gcaggatggg	agaggaagga	cacacttctg	ctcacactgg	ccccaggctc	tagaaciaaag	6000
cagaagcact	cacgggcttg	aattgggtca	ggtggggcct	ccaaggcaa	acccctagcc	6060
caggggtcccc	aacccccaa	tcatagatag	gtactggtcc	atggactggt	aggagccggg	6120
ctgcacagca	cgaagtgagt	ggtgggtgag	tgagcaatac	gcctgagctc	cgcctcctgt	6180
cagatcaagt	ggccttacat	tctcatagga	gtgtgaacct	tattgtgaac	tgacatgcc	6240
agggatctag	gtggagggtc	ccttatgaga	atctaagcc	tgatgatctg	aggtggaaca	6300
gcttcatccc	aaaaccatcc	ccgctggccc	atggaaaaat	tgtccaccac	aaaactgggtc	6360
cctgggtgcca	aaaaggctga	ggaccactgc	tctagccaga	ccttcagaaa	aggaaaatgg	6420
ggccaggcgc	agtggctcgt	gcctgtaatc	ccagcacttt	gggaggccga	ggcaggagga	6480
tcccttgagg	tcaggaattc	aagaccaacc	tggccaacat	ggtgaaacct	catctctgct	6540
aaaaatacaa	aaattagctg	ggtgtggtgg	cgcgtgcctg	taatcccagc	tacttgggag	6600
gctgaggcag	gagaatgggt	tgaacccggg	agacggaggt	tgcaagtgagc	cgagatggca	6660
ccactgcact	ccagcctggg	tgacagaggg	agactccatt	aaaaaaaaaa	gaagaagaaa	6720
gaaagagaga	aaggaagaga	gggaaagaaa	gaaaaaggaa	agaaagaaag	aaagagaaag	6780
aaagaaaaga	aagaaagaaa	ggaaagaaag	aaaggaaaga	aagaaagaaa	gaaagaaaga	6840
aagaaagaaa	gaaagaaaga	aagaaaaata	gaaagagaaa	gaaagggaaa	ggaaaggaaa	6900
atggaaatga	gggctaaaac	ggcgcgggcc	taggactgct	ccgggagaaa	ggagggacct	6960
gggaggaatg	cagtctcaac	tctgtcatct	ctcccgagc	agcctgacaa	aggccctgtg	7020
aggaaaagagg	aggctgaaga	gcacaggccc	aggggaggct	ggagcacctc	cattccattg	7080
ttgggatctc	accagggcag	gggcggtggg	gaatggaggc	agctggcagg	ggggatgggg	7140
agggaggcca	gcgggtgtac	ctggcctgct	gggccacctg	ggactcctgc	acgctgctca	7200
gtgcacctct	ggcgtctctg	gtggcgtgct	tcatgtagcc	ctgcatgaag	ctgagaaggg	7260
aggcatcctc	ggcctctgaa	gctcctgagg	aaagagcagg	gctgagtggg	gtggatcggc	7320
ctctggagca	gccctggggc	tccctgctga	ccaccattg	gactgggagc	cccaagttgc	7380
ctccaccctg	ccccagccc	agtcaccaca	agtgettacg	ggcagaggcc	aggagcgcca	7440
ggagggcaac	aacaaggagt	acccggggct	gcattggacc	tctgttccctg	caaggagtg	7500
tcctgtgagg	ggcaccccag	gtcccatgct	ctctggacct	ctccctgggg	agggtggctg	7560
gccccctaagg	tagaacctta	gctgggtctg	ccagaaggag	taggggccc	ctcctgctca	7620
tacgggctct	cagaaggggg	actggtgagc	ggcgagggat	cgaggcccaa	aggaggtggg	7680
tgggatggag	cagaaaacct	accagactga	acatcaaggc	actgcggtct	ggactgatct	7740
ccgtccagtc	cagccaacat	gctgtgtgtc	tttgggtgat	ttctggccct	cttccaggcc	7800
tcagtttccc	tgtctggggg	aggactgggc	tgtctaaggc	ttcctccagc	cctaagcctg	7860
aagaatgagg	ggggaacctg	cacttgagac	cacttccagc	cccacccctc	atgtagcttt	7920
gggcaagtga	cacctcccgg	gcctccatgt	tcttcagggt	atgatgaggg	gtggggggca	7980
cccgtccagc	tccgaggctt	ccttagctct	agcaagtgct	tctccaggct	tgctggctgg	8040
gctgggcagg	gagctcctct	tgcctctctt	catectctc	ccctcctctt	tccctccccc	8100
agagggcatt	acctggagca	gctgcctcta	gggatgaact	gagcagacag	gcaggagggt	8160
tctgacctgt	tttatatcat	ctccagggca	gcaggcactg	aggaccagg	gcgctgggca	8220
aaggtcacct	gctgaccagt	ggagatgagg	gcctgaggca	gggtgtccag	atgcagcaag	8280
cgggcgggag	agttgggaaa	tccctaggag	actgagtcca	cgctgtctgc	ccggcagccc	8340
tgacagccag	atgagctcag	gaactggggg	tgggcctggg	gagcctcatt	ccttccctagc	8400
tgactggctc	cccagggaga	ggctggtgag	aggggaaatg	gcttggacat	aggccagcgc	8460
tcggggccat	ctcagccttt	cacactggaa	tttcaggccc	ctccctccac	cagccccaag	8520
ccctgaacac	agcctggagt	agaggggtga	ggggcttctt	cagacttgag	aacaagtggg	8580
tggcttgggc	tgggggggtg	ttggagtaaa	ggcacagaag	accgggcac	agtggctcgc	8640
aaggcctggt	tacagggctg	agctcacagc	ccctccagc	acctccatct	ctgggtttca	8700
atccaggcag	gcgagtgtg	ctcaccagc	agccccccac	ccgccccccac	cctgtgtgcc	8760
ccaccgccgc	ctccccctga	gtgtagggca	gggggttggg	gagaagcgca	aggccgctca	8820
gagcccgagg	cctttgcccc	tccctccacc	aggctcccta	ttttgcccct	tggaaccact	8880
gataacatcc	ctgggggagg	agccccacc	aggactgatt	ctctcgttca	ccttccctggg	8940

ttcctcagtg gggctggggg aagagc

8966

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/17908

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A01N 43/04; C07H 21/04; C12N 15/09, 15/11, 15/85

US CL :435/440, 455, 471, 490; 514/44; 536/23.1, 23.5, 24.1, 24.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/440, 455, 471, 490; 514/44; 536/23.1, 23.5, 24.1, 24.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, EMBASE, MEDLINE, DERWENT

search terms: apo? ldl? low density lipoprotein? cholesterol? atherosclerosis? clathrin?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	US 5,736,157 A (WILLIAMS) 07 April 1998, see entire document.	1-4, 7-9, 24-50
A,E	US 4,772,549 A (FROSSARD) 20 September 1988, see entire document.	1-4, 7-9, 24-50

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 DECEMBER 1998

Date of mailing of the international search report

05 FEB 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

NANCY J. DEGEN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/17908

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5-6, 10-23, 51-52
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.